

Pregnancy Risk Assessment Monitoring System

Tennessee
PRAMS
Summary Report
2007



PRAMS

Healthy Babies Start
With Healthy Mothers

Acknowledgments

Tennessee PRAMS project staff would like to express their gratitude to all of the mothers who took time to participate in the survey. Their information will provide a better understanding of the health of mothers and babies in Tennessee.

Project staff would like to acknowledge the PRAMS Steering Committee, as well as Kathy Henson and Colleen Edgehill for their administrative support. We would also like to thank the CDC/PRAMS project manager for Tennessee, LaTreace Q. Harris, for her support and assistance.

Birth certificate data and sampling were provided by the Division of Health Statistics, Tennessee Department of Health.

This publication was made possible by grant number 1 UR6 DP000528 from the Centers for Disease Control and Prevention.

Table of Contents

| | |
|--|-------|
| Executive Summary..... | 1 |
| Overview | |
| Background..... | 2 |
| Methodology..... | 2-3 |
| Technical Notes..... | 3 |
| PRAMS Staff and Steering Committee..... | 4 |
| Results | |
| Pregnancy Intention..... | 6-10 |
| Folic Acid Awareness and Multivitamin Use..... | 11-14 |
| Prenatal Care..... | 15-20 |
| HIV Discussion and Testing..... | 21-24 |
| Physical Abuse..... | 25-27 |
| Stress..... | 28-32 |
| Cigarette Smoking..... | 33-36 |
| Alcohol Use..... | 37-39 |
| Maternal Infections..... | 40-42 |
| Maternal Health Problems..... | 43-45 |
| WIC Participation..... | 46-48 |
| Breastfeeding..... | 50-52 |
| Dental Care..... | 53-55 |
| Postpartum Depression..... | 56-58 |
| Postpartum Birth Control..... | 59-61 |
| Postpartum Checkup..... | 62-63 |
| Infant Health Care..... | 64-66 |
| Sleep Behaviors..... | 67-69 |
| Infant Smoke Exposure..... | 70-71 |
| Infant Safety..... | 72-73 |
| Low Birthweight..... | 74-75 |
| Appendix – 2007 TN PRAMS Questionnaire | 76-91 |

Executive Summary

The Tennessee Pregnancy Risk Assessment Monitoring System (TN PRAMS) is a population-based survey of randomly sampled women who gave birth to a live-born infant in Tennessee. The topics included in the survey were selected based on their relevance to maternal and infant health. The following summary highlights important findings within the report:

- One-half (51.5%) of pregnancies were unintended.
- Among women *not* trying to get pregnant, approximately one-half were also not using birth control.
- Three-fourths (73.0%) of mothers did not take a daily multivitamin in the month prior to pregnancy.
- One-fifth (21.5%) of women received late or no prenatal care.
- Approximately 43% of women who received prenatal care reported that it was either partially or totally paid for by TennCare.
- Among women who received prenatal care, 68.2% reported discussing HIV testing and 69.7% reported having an HIV test.
- Approximately 11% of women reported physical abuse before and/or during pregnancy.
- A majority of women (77.9%) reported at least one stressor in the 12 months prior to delivery, with financial-related problems being the most frequently reported type of stressor.
- One-fifth (21.2%) of women smoked cigarettes and 5.8% drank alcohol during the last 3 months of pregnancy.
- Two-fifths of women (40.2%) were diagnosed with a urinary tract infection, sexually transmitted disease or other vaginal infection during pregnancy.
- One-half (50.9%) of women were enrolled in the Supplemental Nutrition Program for Women, Infants and Children (WIC) during pregnancy.
- One-third (31.4%) of mothers did not initiate breastfeeding, with not liking breastfeeding being the most frequently reported reason for not doing so.
- Overall, approximately two-thirds (66.6%) of women did not see a dentist during pregnancy. Among those with a reported dental problem, over one-half did not go to a dentist or dental clinic.
- Approximately 20% of women reported symptoms of postpartum depression.
- The majority of mothers (89.2%) were using postpartum birth control at the time of the survey.
- The majority of mothers (87.2%) had received a postpartum checkup for themselves, and 97.5% reported that their infant had had a well-baby checkup.
- One-third (37.5%) of mothers most often placed their infant to sleep on his or her stomach or side, and one-half (50.9%) reported bed sharing with their infant.
- Approximately 9% of mothers reported that their infants were exposed to secondhand smoke.
- Almost all mothers reported using a car seat to bring their infant home from the hospital (99.2%) and always or almost always using a car seat (99.7%).
- Approximately 5% of women reported that there is not a working smoke alarm in their home, and 8% reported that there are loaded firearms in their home.
- Approximately 8.5% of mothers gave birth to low birthweight infants.

Overview

Background

Infant mortality rates in Tennessee have remained unchanged in recent years and have been consistently higher than national rates. Racial disparities have also persisted, with black babies two-and-a-half times as likely as white babies to die during their first year of life. Racial disparities in infant mortality are related in-part to differences in recognized risk factors for infant mortality. Based on birth certificate data in Tennessee, the prevalence of the following risk factors are all higher among infants born to black mothers than among those born to white mothers: low birthweight, premature birth, multiple birth, teenage mother, unmarried mother, mother with a previous child death, mother with high school or lower education, and no prenatal care. In order to reduce infant mortality in Tennessee, these risk factors will need to be addressed. However, the above information is limited to data collected on birth and death certificates and to a large extent the risk factors listed occur *during* pregnancy. While this information is vital to understanding and reducing the burden of infant mortality in the state, it doesn't address risk factors that occur *prior to* or *after* pregnancy.

The Tennessee Department of Health (TDH) recognizes the need for maternal and child health data beyond that available from our current systems of vital and programmatic records. The Tennessee Pregnancy Risk Assessment Monitoring System (TN PRAMS), which provides information on maternal behaviors and experiences prior to, during *and* after pregnancy, will provide data vital to our efforts to target programs and activities appropriately, and to ultimately improve birth outcomes and the health of both mothers and children in the state.

Methodology

Tennessee PRAMS is conducted by TDH, Office of Policy, Planning and Assessment (PPA), Research Section. The project was established in 2006 through a collaborative agreement between the Centers for Disease Control and Prevention (CDC) and TDH. Tennessee PRAMS was designed to collect, analyze, and disseminate information on a variety of maternal behaviors and experiences that may be associated with various birth outcomes.

Tennessee PRAMS is a population-based survey of new mothers. In 2007, approximately 1,200 mothers were randomly selected from the Tennessee Vital Statistics birth file to participate in the survey. In order to be eligible for selection, women had to be Tennessee residents and have delivered a live-born infant. Women were selected using stratified random sampling. Mothers were first separated into two different groups (or strata) based on infant birthweight and each group was then randomly sampled. Women in the low birthweight (LBW) group were sampled at a higher rate than those in the normal birthweight (NBW) group (i.e. women giving birth to LBW infants were oversampled). Oversampling makes it possible to better estimate risks for groups that are relatively small compared to the rest of the population.

Selected mothers were first notified of the PRAMS survey and then sent the questionnaire via mail approximately 2-6 months after delivery. If the mother did not respond after three attempts by mail, she was then contacted by telephone and given the opportunity to complete the questionnaire via phone interview. The questionnaire consisted of 83 standardized questions (see Appendix). There were 4 versions of the questionnaire available: English adult, Spanish adult, English teen and Spanish teen. Mothers whose ethnicity was marked as 'Hispanic' on the birth certificate were sent both an English and Spanish version. Mothers under 18 years of age were sent the teen version(s). A bilingual telephone interviewer was available for mothers more comfortable communicating in Spanish.

Overview cont.

After data collection was concluded, mothers' responses were linked to their corresponding birth certificate data. This linked PRAMS response/birth certificate dataset was then sent to the CDC for weighting. Weighting allows for estimation of statistics for the entire state's population of women who delivered a live-born infant based on data gathered from a small sample of mothers in that population. Tennessee PRAMS data is weighted based on sample design, non-response and non-coverage. In 2007, 705 out of 1,160 sampled mothers completed the questionnaire and the overall, weighted response rate was 63%. The weighted response rates among mothers of LBW and NBW infants were 59% and 64%, respectively. The CDC regards 70% to be the minimum acceptable response rate and does not publish data from states with response rates below this threshold. ***Because the 2007 TN PRAMS did not achieve a 70% response rate, results obtained using this data may not be statistically valid and must be interpreted with caution. It is recommended that the information in this report be maintained for internal Health Department use only.***

Technical Notes

All data in this report were analyzed using the proc surveyfreq procedure in SAS 9.2. Most data are presented as simple descriptive statistics and two-way comparisons. Each topic area is organized to include tables which present the prevalence of indicators by select demographic and socioeconomic characteristics (race/ethnicity, age, marital status, education, income, insurance), as well as graphical presentations of subgroup analyses. All prevalence tables include 95% confidence intervals, as do select graphs. Two-way comparisons for subgroup analyses were tested for statistical significance using a chi-square test, with p-values of less than 0.05 considered significant. Unless otherwise indicated, differences noted in the text are statistically significant.

Because estimates based on small samples are imprecise and may be biased, estimates for which the number of respondents was fewer than 30 (unweighted sample size) are not reported. Estimates based on sample sizes between 30 and 60 are reported, but they include a note stating that the results may be unreliable. Hispanic women are included in most prevalence tables. However, it is important to note that only 42 Hispanics completed the survey and these prevalence rates should therefore be interpreted with caution. For some indicators there were fewer than 30 Hispanic respondents, and in these instances data for Hispanic mothers are excluded from the prevalence tables.

Insurance status in all prevalence tables was defined based on women's insurance status just prior to becoming pregnant. Women who reported being on Medicaid, TennCare or CoverKids (with or without other health insurance) were classified in the Medicaid group. Women who reported having insurance but not being on Medicaid, TennCare or CoverKids were classified in the insurance group.

Overview *cont.*

PRAMS Staff and Steering Committee

The TN PRAMS Project staff for 2007 consisted of:

- David J. Law, PhD (PRAMS Project Director)
- Anthony H. Rico, MS (PRAMS Project Coordinator)
- Audrey M. Bauer, DVM, MPH (Epidemiologist/Analyst)
- Yinmei Li, MD, MPH (Epidemiologist)
- David Howard (PRAMS Data Manager)

The TN PRAMS Steering Committee for 2007 advised staff on the development and selection of state-specific questions and on the use, dissemination, and application of findings. Committee members included:

- David J. Law, PhD, Director of Research, PPA, TDH
- Sharon Leinbach, State Registrar, TDH
- Tom Spillman, Statistical Analyst Supervisor, TDH
- Theodora Pinnock, MD, Director of Maternal and Child Health, TDH
- Paula Taylor, MS, Director, PPA, TDH
- Margaret Major, RD, MPA, Chief, Women's and Genetics Section, TDH
- Peggy Lewis, Director of WIC, TDH
- Robbie Jackman, Director of Minority Health, TDH
- Cheryl W. Major, RNC, BSN, President, Tennessee Perinatal Association
- Mark Gaylord, MD, University of Tennessee-Knoxville and Tennessee Pediatric Society
- Melanie Lutenbacher, PhD, APRN, Director of Nursing Science Programs, Vanderbilt University
- Cathy R. Taylor, MSN, RN, DrPH, Assistant Professor, Meharry/Vanderbilt Alliance
- Kimberlee Wyche-Etheridge, MD, MPH, Maternal Child Health Director, Nashville-Davidson County Metropolitan Health Department

2007
Tennessee
PRAMS
Results

Pregnancy Intention

Information regarding pregnancy intention was derived from the following questions:

Question #12: Thinking back to just before you got pregnant, how did you feel about becoming pregnant?

- I wanted to be pregnant sooner
- I wanted to be pregnant later
- I wanted to be pregnant then
- I didn't want to be pregnant then or at any time in the future

Question #14: When you got pregnant with your new baby, were you trying to get pregnant?

- No
- Yes


An intended pregnancy was one in which the mother answered that she wanted to be pregnant then or sooner. Women who wanted to be pregnant later or not at all were classified as having an unintended pregnancy. Unintended pregnancies were further subdivided into two categories: mistimed pregnancies or unwanted pregnancies. Mistimed pregnancies were those in which the mother wanted to be pregnant later than the time she became pregnant. Unwanted pregnancies were those in which the mother did not want to be pregnant then or anytime in the future.

Additional questions addressing birth control use at the time of pregnancy (Questions #15-17) may be found in the Appendix.

Key Findings:

- Approximately one-half (51.5%) of pregnancies were unintended.
- Among unintended pregnancies, 74.3% were mistimed and 25.7% were unwanted.
- Unintended pregnancies were more common among black non-Hispanics than among white non-Hispanics.
- There were more unintended pregnancies among unmarried women than among married women.
- Unintended pregnancies decreased with increasing age, education and income.
- Among women who were *not* trying to get pregnant, approximately one-half were using birth control and one-half were not using birth control.
 - The percentage of women not using birth control was similar across demographic and socioeconomic subgroups.
 - Among women who were not using birth control, the most frequently reported reason for not doing so was not minding getting pregnant.
 - Among women who were using birth control, the most frequently used method at the time of pregnancy was condoms.
- Compared to women with intended pregnancies, those with unintended pregnancies were more likely to:
 - receive no preconception care,
 - receive late or no prenatal care, and
 - not take a daily multivitamin in the month prior to pregnancy.
- However, even among women with *intended* pregnancies, approximately two-thirds did not receive preconception care or take a daily multivitamin in the month prior to pregnancy.

Pregnancy Intention *cont.*




Unintended Pregnancies by Demographic and Socioeconomic Characteristics Tennessee, 2007

| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 45.9 | 40.1-59.9 |
| | Black Non-Hisp | 73.3 | 62.5-84.1 |
| | Hispanic | 45.8 | 26.2-65.4 |
| Age (years) | <20 | 75.7 | 64.0-87.4 |
| | 20-29 | 52.7 | 46.0-59.4 |
| | 30+ | 36.6 | 27.9-45.2 |
| Married | Yes | 35.1 | 29.2-41.1 |
| | No | 72.4 | 65.0-79.7 |
| Education (years) | <12 | 62.7 | 52.3-73.0 |
| | 12 | 54.5 | 44.5-64.4 |
| | >12 | 43.3 | 36.5-50.2 |
| Household Income | <\$10,000 | 73.6 | 64.6-82.5 |
| | \$10-24,999 | 59.0 | 46.0-72.1 |
| | \$25-34,999 | 49.2 | 37.4-61.0 |
| | \$35,000+ | 30.4 | 23.0-37.7 |
| Insurance | Medicaid | 72.6 | 62.9-82.3 |
| | Insurance | 35.6 | 28.7-42.4 |

Pregnancy Intention *cont.*

**Birth Control Use Among Women Not Trying to Get Pregnant *
Tennessee, 2007**




**Birth Control Use by Demographic and Socioeconomic Characteristics*
Tennessee, 2007**

| | | Percent <i>Not</i> Using Birth Control | 95% Confidence Interval |
|--------------------------|-----------------------|--|-------------------------|
| Race | White Non-Hisp | 50.3 | 42.3-58.4 |
| | Black Non-Hisp | 46.6 | 31.7-61.5 |
| Age (years) | <20 | 48.2 | 31.0-65.4 |
| | 20-29 | 46.9 | 37.8-55.9 |
| | 30+ | 52.0 | 38.7-65.3 |
| Married | Yes | 52.6 | 43.1-62.1 |
| | No | 45.1 | 35.4-54.8 |
| Education (years) | <12 | 48.4 | 35.1-61.7 |
| | 12 | 52.4 | 39.1-65.6 |
| | >12 | 44.5 | 34.6-54.5 |
| Household Income | <\$10,000 | 47.8 | 34.9-60.6 |
| | \$10-24,999 | 50.8 | 34.3-67.2 |
| | \$25-34,999 | 39.4 | 24.9-54.0 |
| | \$35,000+ | 49.0 | 36.0-62.0 |
| Insurance | Medicaid | 54.9 | 41.6-68.3 |
| | Insurance | 45.8 | 35.1-56.5 |


* Analysis limited to women not trying to get pregnant, regardless of pregnancy intention.

Pregnancy Intention *cont.*

Reason for Not Using Birth Control*
Tennessee, 2007




Birth Control Method at Time of Pregnancy*
Tennessee, 2007




* Women may have reported more than one reason for not using birth control or more than one birth control method. Therefore, percentages do not sum to 100%.

Pregnancy Intention *cont.*


No Preconception Care by Pregnancy Intent
Tennessee, 2007



Late or No Prenatal Care by Pregnancy Intent
Tennessee, 2007



No Daily Multivitamin in Month Prior to Pregnancy by Pregnancy Intent
Tennessee, 2007



Folic Acid Awareness and Multivitamin Use

Information regarding folic acid awareness and multivitamin use was derived from the following questions:

Question #26: Have you ever heard or read that taking the vitamin folic acid can help prevent some birth defects?
_No
_Yes


Question #27: Have you ever heard about folic acid from any of the following?
_Magazine or newspaper article
_Radio or television
_Doctor, nurse or other health care worker
_Book
_Family or friends
_Other

Question #3: During the month before you got pregnant with your new baby, how many times a week did you take a multivitamin or a prenatal vitamin?
_I didn't take a multivitamin or a prenatal vitamin at all
_1 to 3 times per week
_4 to 6 times per week
_Every day of the week

Key Findings:

- Approximately one-fifth (19.0%) of mothers were unaware of the benefits of folic acid.
- Teenage mothers were more likely than older mothers to be unaware of the benefits of folic acid.
- Among women who were aware of the benefits of folic acid, the most frequently reported source of information was a doctor, nurse or other health care worker.
- In addition to the sources of folic acid information listed in Question #27, other reported sources included the internet and through school and/or work (e.g. nursing, pharmacy).
- Approximately three-fourths (73.0%) of mothers did not take a daily multivitamin in the month prior to pregnancy.
- There was no statistically significant difference in daily multivitamin use among women who were versus those who were not aware of the benefits of folic acid. However, it was not possible to determine *when* women became aware of folic acid (i.e. before, during or after pregnancy).
- Compared to women with intended pregnancies, those with unintended pregnancies were more likely to not take a daily multivitamin (see page 10).
- There was no statistically significant difference in the percentage of women who were unaware of the benefits of folic acid among those with versus those without preconception care. However, women who received preconception care were more likely than those who did not receive care to take a daily multivitamin.

Folic Acid Awareness and Multivitamin Use *cont.*




**Folic Acid Awareness by Demographic and Socioeconomic Characteristics
Tennessee, 2007**

| | | Percent <i>Unaware</i> of Folic Acid | 95% Confidence Interval |
|--------------------------|-----------------------|---|-------------------------|
| Race | White Non-Hisp | 18.3 | 13.7-22.8 |
| | Black Non-Hisp | 25.4 | 14.2-36.6 |
| | Hispanic | 14.1 | 0.0-28.4 |
| Age (years) | <20 | 39.4 | 25.2-53.6 |
| | 20-29 | 18.5 | 13.2-23.8 |
| | 30+ | 10.0 | 4.6-15.4 |
| Married | Yes | 12.0 | 7.9-16.1 |
| | No | 27.8 | 20.4-35.2 |
| Education (years) | <12 | 28.1 | 18.4-37.9 |
| | 12 | 23.4 | 14.8-32.0 |
| | >12 | 11.9 | 7.3-16.5 |
| Household Income | <\$10,000 | 25.2 | 16.0-34.4 |
| | \$10-24,999 | 23.6 | 12.5-34.8 |
| | \$25-34,999 | 20.5 | 10.9-30.1 |
| | \$35,000+ | 8.3 | 3.9-12.8 |
| Insurance | Medicaid | 31.0 | 20.9-41.1 |
| | Insurance | 9.8 | 5.6-14.1 |

Folic Acid Awareness and Multivitamin Use *cont.*

Frequency of Multivitamin Use In Month Prior to Pregnancy
Tennessee, 2007




No Daily Multivitamin by Demographic and Socioeconomic Characteristics
Tennessee, 2007


| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 74.2 | 69.3-79.1 |
| | Black Non-Hisp | 66.9 | 55.4-78.4 |
| | Hispanic | 80.6 | 64.7-96.5 |
| Age (years) | <20 | 79.8 | 68.2-91.5 |
| | 20-29 | 72.6 | 66.7-78.4 |
| | 30+ | 70.6 | 62.8-78.4 |
| Married | Yes | 67.2 | 61.4-72.9 |
| | No | 80.5 | 73.9-87.0 |
| Education (years) | <12 | 71.8 | 62.1-81.6 |
| | 12 | 78.8 | 70.9-86.7 |
| | >12 | 71.0 | 65.0-76.9 |
| Household Income | <\$10,000 | 74.5 | 65.4-83.5 |
| | \$10-24,999 | 80.0 | 69.7-90.3 |
| | \$25-34,999 | 82.0 | 73.1-90.9 |
| | \$35,000+ | 64.8 | 57.3-72.2 |
| Insurance | Medicaid | 75.5 | 66.3-84.6 |
| | Insurance | 69.0 | 62.7-75.2 |

Folic Acid Awareness and Multivitamin Use *cont.*


**Sources of Folic Acid Information
Tennessee, 2007***




**Number of Sources of Folic Acid Information
Tennessee, 2007**



**No Daily Multivitamin Use by Folic Acid Awareness
Tennessee, 2007†**



**Folic Acid Awareness and Multivitamin Use by Preconception Care
Tennessee, 2007‡**



* Women may have reported more than one source of folic acid information. Therefore, percentages do not sum to 100%.

† Not statistically significant.

‡ Analysis limited to women with intended pregnancies.

Prenatal Care

Information regarding the timing of prenatal care was derived from the following question:

Question #19: How many weeks or months pregnant were you when you had your first visit for prenatal care?

[#] Weeks

[#] Months

_I did not go for prenatal care


Mothers who initiated care after the first trimester of pregnancy (after 12 weeks/3 months or later) were classified as receiving late prenatal care. Those who initiated care within the first trimester were classified as receiving early prenatal care.

Additional questions addressing the desired timing of prenatal care (Question #20), barriers to getting prenatal care (Question #21), payment source (Question #22) and prenatal care content (Question #23) may be found in the Appendix.

Key Findings:

- Approximately one-fifth (21.5%) of women received late or no prenatal care.
- Black non-Hispanics were more likely than white non-Hispanics to receive late or no care.
- Unmarried women were more likely than married women to receive late or no care.
- The prevalence of late or no prenatal care increased with decreasing age, education and income.
- Women with unintended pregnancies were more likely to receive late or no care than those with intended pregnancies (see pg 10).
- Women who recognized their pregnancy late (after the first 8 weeks of gestation) were more likely to receive late or no care than those who recognized their pregnancy earlier.
- Approximately one-fifth (18.5%) of women did not receive prenatal care as early as wanted.
- Approximately two-thirds of women who received late prenatal care did not get care as early as wanted.
- Approximately one-third (32.8%) of women reported at least one barrier to care.
- The most frequently reported barriers to receiving prenatal care were lack of money/insurance and not being able to get an appointment.
- The percentage of women reporting at least one barrier to care was higher among those who received late care than among those who received early care.
- Almost one-half of women reported that their prenatal care was paid for partially or completely by TennCare.
- The most frequently discussed topics during prenatal care were medicine safety, birth defects screening, early labor and breastfeeding. The least frequently discussed topics were seat belt use and physical abuse.

Prenatal Care *cont.*




**Late or No Prenatal Care by Demographic and Socioeconomic Characteristics
Tennessee, 2007**


| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 17.1 | 12.4-21.8 |
| | Black Non-Hisp | 32.5 | 20.3-44.7 |
| | Hispanic | 39.1 | 18.5-59.7 |
| Age (years) | <20 | 33.6 | 18.9-48.3 |
| | 20-29 | 23.2 | 17.2-29.2 |
| | 30+ | 12.5 | 6.1-18.8 |
| Married | Yes | 12.9 | 8.6-17.2 |
| | No | 32.6 | 24.4-40.8 |
| Education (years) | <12 | 32.9 | 22.0-43.7 |
| | 12 | 20.4 | 12.0-28.8 |
| | >12 | 16.7 | 11.0-22.3 |
| Household Income | <\$10,000 | 44.6 | 33.5-55.8 |
| | \$10-24,999 | 24.5 | 13.0-36.0 |
| | \$25-34,999 | 15.4 | 6.7-24.1 |
| | \$35,000+ | 4.7 | 1.5-7.8 |
| Insurance | Medicaid | 29.7 | 19.0-40.4 |
| | Insurance | 6.3 | 2.6-9.9 |

Prenatal Care *cont.*

Received Prenatal Care as Early as Wanted*
Tennessee, 2007



Did Not Receive Prenatal Care as Early as Wanted by Timing of Care
Tennessee, 2007




**Did Not Receive Care as Early as Wanted
by Demographic and Socioeconomic Characteristics*
Tennessee, 2007**

| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 15.9 | 11.3-20.6 |
| | Black Non-Hisp | 31.1 | 18.7-43.6 |
| | Hispanic | 16.6 | 0.7-32.5 |
| Age (years) | <20 | 29.7 | 15.1-44.4 |
| | 20-29 | 21.4 | 15.3-27.4 |
| | 30+ | 7.9 | 3.1-12.8 |
| Married | Yes | 9.8 | 6.0-13.7 |
| | No | 30.0 | 21.9-38.2 |
| Education (years) | <12 | 22.0 | 12.1-31.9 |
| | 12 | 19.6 | 11.1-28.1 |
| | >12 | 16.4 | 10.7-22.2 |
| Household Income | <\$10,000 | 35.4 | 24.3-46.4 |
| | \$10-24,999 | 26.6 | 14.3-38.9 |
| | \$25-34,999 | 17.9 | 8.4-27.4 |
| | \$35,000+ | 3.4 | 0.9-5.8 |
| Insurance | Medicaid | 25.4 | 15.0-35.7 |
| | Insurance | 9.7 | 5.1-14.4 |


* Analysis limited to women who received prenatal care.

Prenatal Care *cont.*

Barriers to Prenatal Care^{†§}
Tennessee, 2007



Barriers to Prenatal Care by Timing of Care^{‡§}
Tennessee, 2007



* Not statistically significant.


† Analysis limited to women who received prenatal care.

‡ Women may have reported more than one barrier to care. Therefore, percentages do not sum to 100%.


§ Other barriers to care included "didn't know/believe I was pregnant" and "took too long to get coverage", "did not know where to go" or other utilization/availability issues.

Prenatal Care *cont.*

Number of Barriers to Prenatal Care*
Tennessee, 2007



Number of Barriers to Prenatal Care by Timing of Care
Tennessee, 2007




One or More Barriers to Prenatal Care by Demographic and Socioeconomic Characteristics* Tennessee, 2007

| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 31.0 | 25.0-36.9 |
| | Black Non-Hisp | 33.8 | 21.2-46.3 |
| | Hispanic | 52.7 | 31.8-73.5 |
| Age (years) | <20 | 35.9 | 20.2-51.7 |
| | 20-29 | 36.1 | 29.2-43.1 |
| | 30+ | 24.7 | 16.5-33.0 |
| Married | Yes | 25.7 | 19.9-31.6 |
| | No | 41.8 | 33.0-50.6 |
| Education (years) | <12 | 42.1 | 30.5-53.7 |
| | 12 | 35.0 | 24.6-45.4 |
| | >12 | 27.4 | 20.7-34.1 |
| Household Income | <\$10,000 | 44.4 | 32.6-56.2 |
| | \$10-24,999 | 50.6 | 37.0-64.2 |
| | \$25-34,999 | 31.2 | 19.8-42.6 |
| | \$35,000+ | 15.1 | 9.0-21.2 |
| Insurance | Medicaid | 33.3 | 21.9-44.6 |
| | Insurance | 17.5 | 11.6-23.5 |


* Analysis limited to women who received prenatal care.

Prenatal Care *cont.*


**Prenatal Care Payment Source*
Tennessee, 2007**



**Number of Prenatal Care Payment Sources
Tennessee, 2007**



**Prenatal Care Content*
Tennessee, 2007**



* Women may have reported more than one source of payment or prenatal care topic. Therefore, percentages do not sum to 100%.

HIV Discussion and Testing

Information regarding HIV discussion and testing was derived from the following questions:

Question #23j: During any of your prenatal care visits, did a doctor, nurse or other health care worker talk with you about getting tested for HIV (the virus that causes AIDS)?

- _No
- _Yes

Question #24: At any time during your most recent pregnancy or delivery, did you have a test for HIV?

- _No
- _Yes
- _I don't know

Question #25: Were you offered an HIV test during your most recent pregnancy or delivery?

- _No
- _Yes


All analyses of HIV discussion and testing were limited to women who received prenatal care.

Key Findings:

- Among women who received prenatal care, 68.2% reported discussing HIV testing and 69.7% reported having an HIV test.
- Black non-Hispanics were more likely than white non-Hispanics to report discussion of HIV testing during prenatal care and to have received an HIV test.
- There was no statistically significant difference in the percentage of women who had an HIV test among those who received prenatal care early versus those who received care late.
- Women whose prenatal care included a discussion of HIV testing were more likely to have received an HIV test than women whose prenatal care did not include such a discussion.
- The majority of women (88.5%) who did not have an HIV test reported that they were not offered an HIV test. It was not possible to determine why the remaining 11.5% of women who were offered an HIV test did not receive one.

HIV Discussion and Testing *cont.*

Discussion of HIV Testing During Prenatal Care
Tennessee, 2007




HIV Discussion by Demographic and Socioeconomic Characteristics
Tennessee, 2007

| | | Percent | 95% Confidence Interval |
|-------------------|----------------|---------|-------------------------|
| Race | White Non-Hisp | 62.7 | 56.9-68.5 |
| | Black Non-Hisp | 83.0 | 73.4-92.6 |
| | Hispanic | 85.5 | 72.3-98.8 |
| Age (years) | <20 | 69.5 | 55.0-84.0 |
| | 20-29 | 66.4 | 59.9-73.0 |
| | 30+ | 71.1 | 63.0-79.1 |
| Married | Yes | 65.2 | 59.1-71.2 |
| | No | 72.2 | 64.4-80.1 |
| Education (years) | <12 | 80.3 | 71.2-89.4 |
| | 12 | 72.3 | 62.7-81.9 |
| | >12 | 60.1 | 53.3-66.9 |
| Household Income | <\$10,000 | 75.2 | 65.2-85.2 |
| | \$10-24,999 | 64.9 | 51.8-77.9 |
| | \$25-34,999 | 73.0 | 62.6-83.4 |
| | \$35,000+ | 60.2 | 52.3-68.2 |
| Insurance | Medicaid | 71.1 | 60.6-81.7 |
| | Insurance | 65.2 | 58.4-72.0 |

HIV Discussion and Testing *cont.*

HIV Testing During Pregnancy or Delivery
Tennessee, 2007




HIV Testing by Demographic and Socioeconomic Characteristics
Tennessee, 2007


| | | Percent | 95% Confidence Interval |
|-------------------|----------------|---------|-------------------------|
| Race | White Non-Hisp | 66.6 | 60.6-72.7 |
| | Black Non-Hisp | 82.4 | 71.6-93.1 |
| | Hispanic | 66.3 | 45.2-87.4 |
| Age (years) | <20 | 69.8 | 54.4-85.1 |
| | 20-29 | 71.5 | 64.8-78.2 |
| | 30+ | 66.5 | 57.3-75.6 |
| Married | Yes | 64.7 | 58.2-71.1 |
| | No | 76.8 | 68.7-84.9 |
| Education (years) | <12 | 70.4 | 59.0-81.7 |
| | 12 | 73.9 | 63.7-84.1 |
| | >12 | 66.9 | 60.0-73.8 |
| Household Income | <\$10,000 | 80.8 | 70.8-90.8 |
| | \$10-24,999 | 69.4 | 56.1-82.8 |
| | \$25-34,999 | 72.2 | 60.6-83.8 |
| | \$35,000+ | 62.3 | 53.9-70.7 |
| Insurance | Medicaid | 77.0 | 66.3-87.8 |
| | Insurance | 65.2 | 57.9-72.5 |

HIV Discussion and Testing *cont.*


**HIV Testing by Timing of Prenatal Care*
Tennessee, 2007**



**HIV Testing by Prenatal Care Content
Tennessee, 2007**



**Offer of HIV Test Among Women Without a Test
Tennessee, 2007**



* Not statistically significant.

Physical Abuse

Information regarding physical abuse was derived from the following questions:

Question #42a: During the 12 months before you got pregnant, did an ex-husband or ex-partner push, hit, slap, kick, choke, or physically hurt you in any other way?

- _No
- _Yes

Question #42b: During the 12 months before you got pregnant, were you physically hurt in any way by your husband or partner?

- _No
- _Yes

Additional (and similarly worded) questions about abuse *during* pregnancy (Questions #43a and #43b) may be found in the Appendix. Except where otherwise noted, a woman was considered to be physically abused if she responded positively to any of the four physical abuse questions noted above. In other words, physical abuse was defined as having been pushed, hit, slapped, kicked, choked or otherwise hurt by a husband/partner and/or ex-husband/partner prior to and/or during pregnancy.


Due to reporting requirements in Tennessee, mothers who were under 18 years of age were excluded from answering physical abuse questions and are therefore not included in the analysis.

Key Findings:


- Approximately 11% of women reported physical abuse before and/or during pregnancy.
 - Among women who were abused, 42% reported abuse prior to pregnancy, 18% reported abuse during pregnancy and 40% reported abuse both prior to *and* during pregnancy.
 - Among women who were abused, 12% reported abuse by their husband/partner, 32% reported abuse by their ex-husband/partner and 56% reported abuse by both their current *and* ex-husband/partner.
- Women less than 30 years of age were more likely than older women to report physical abuse.
- Unmarried women were more likely than those who were married to report physical abuse.
- Compared to women who were not abused, those who reported physical abuse were more likely to:
 - Have an unintended pregnancy,
 - Receive late or no prenatal care,
 - Experience one or more stressful events in the 12 months prior to delivery,
 - Be diagnosed with an infection during pregnancy (see page 42), and
 - Report symptoms of postpartum depression.
- Women who were physically abused were less likely to initiate breastfeeding than women who did not report abuse.

Physical Abuse cont.

Timing of Physical Abuse
Tennessee, 2007



Physical Abuse by Abuser
Tennessee, 2007




Physical Abuse by Demographic and Socioeconomic Characteristics Tennessee, 2007

| | | Percent | 95% Confidence Interval |
|--------------------------|----------------|---------|-------------------------|
| Race | White Non-Hisp | 10.7 | 6.8-14.6 |
| | Black Non-Hisp | 15.2 | 5.6-24.9 |
| | Hispanic | 3.7 | 0.0-10.9 |
| Age (years) | 19* | 26.4 | 10.0-42.8 |
| | 20-29 | 12.6 | 7.9-17.3 |
| | 30+ | 2.5 | 0.0-5.1 |
| Married | Yes | 2.5 | 0.6-4.4 |
| | No | 22.6 | 15.3-30.0 |
| Education (years) | <12 | 13.9 | 5.4-22.3 |
| | 12 | 14.5 | 7.0-22.1 |
| | >12 | 6.8 | 3.1-10.5 |
| Household Income | <\$10,000 | 18.0 | 9.2-26.8 |
| | \$10-24,999 | 15.1 | 4.9-25.4 |
| | \$25-34,999 | 16.1 | 7.2-24.9 |
| | \$35,000+ | 0.5 | 0.1-0.9 |
| Insurance | Medicaid | 28.5 | 17.7-39.2 |
| | Insurance | 3.9 | 1.0-6.8 |


* Results for 19 year olds are based on an unweighted sample size of 48. Therefore, results may be unreliable and should be interpreted with caution.

Physical Abuse cont.


**Unintended Pregnancies by Physical Abuse
Tennessee, 2007**




**Late or No Prenatal Care by Physical Abuse*
Tennessee, 2007**




**One or More Stressors by Physical Abuse
Tennessee, 2007**



**Postpartum Depression by Physical Abuse
Tennessee, 2007**



**Breastfeeding by Physical Abuse
Tennessee, 2007**



* Results for abused women are based on an unweighted sample size of 57. Therefore, results may be unreliable and should be interpreted with caution.

Stress

Information regarding stress was derived from the following question:

Question #38: This question is about things that may have happened during the 12 months before your new baby was born. For each item, circle Y (yes) if it happened to you or circle N (no) if it did not.

- a. A close family member was very sick and had to go into the hospital
- b. I got separated or divorced from my husband or partner
- c. I moved to a new address
- d. I was homeless
- e. My husband or partner lost his job
- f. I lost my job even though I wanted to go on working
- g. I argued with my husband or partner more than usual
- h. My husband or partner said he didn't want me to be pregnant
- i. I had a lot of bills I couldn't pay
- j. I was in a physical fight
- k. I or my husband or partner went to jail
- l. Someone very close to me had a bad problem with drinking or drugs
- m. Someone very close to me died


Mothers who responded negatively to all of the above items were classified as having no stress during the 12 months prior to delivery. Those who responded positively to 1, 2 or 3 of the above items were classified as having low stress, and those who responded positively to 4 or more items were classified as having high stress. In addition, the above items were grouped by type of stressor. Women were classified as having emotional stressors (responded positively to items a and/or m), financial stressors (c, e, f and/or i), partner stressors (b, g and/or h) and/or traumatic stressors (d, j, k and/or l).

Additional questions addressing food security (Question #39), neighborhood safety (Question #40) and racial bias (Question #41) may be found in the Appendix.

Key Findings:

- A majority of women (77.9%) reported at least one stressor in the 12 months prior to delivery.
- On average, women reported 2.5 stressors in the 12 months prior to delivery.
- The most frequently reported individual stressor was moving to a new address.
- Financial-related problems were the most frequently reported type of stressor.
- Among women who reported at least one stressor, approximately one-third experienced one type of stressor, while 10% experienced all four types of stressors.
- Among women who reported at least one stressor, approximately one-third experienced high stress levels and two-thirds experienced low stress levels.
- Approximately 14% of women reported that during the 12 months prior to delivery they ate less than they felt they should because there wasn't enough money to buy food (i.e. were food insecure).
- Approximately 13% of women reported that during the 12 months prior to delivery they always, often or sometimes felt unsafe in the neighborhood where they lived.
- Approximately 10% of women reported that during the 12 months prior to delivery they felt emotionally upset as a result of how they were treated based on their race.
- Black non-Hispanics were more likely than white non-Hispanics to report racial bias.

Stress cont.




One or More Stressors by Demographic and Socioeconomic Characteristics Tennessee, 2007


| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 77.8 | 73.3-82.4 |
| | Black Non-Hisp | 87.0 | 79.3-94.8 |
| | Hispanic | 62.9 | 43.7-82.2 |
| Age (years) | <20 | 84.7 | 74.0-95.4 |
| | 20-29 | 82.6 | 77.8-87.4 |
| | 30+ | 64.9 | 56.7-73.1 |
| Married | Yes | 68.7 | 63.1-74.4 |
| | No | 89.3 | 84.3-94.3 |
| Education (years) | <12 | 78.3 | 69.4-87.2 |
| | 12 | 87.9 | 82.0-93.8 |
| | >12 | 71.8 | 65.9-77.7 |
| Household Income | <\$10,000 | 87.9 | 81.2-94.6 |
| | \$10-24,999 | 88.4 | 80.2-96.6 |
| | \$25-34,999 | 88.6 | 81.9-95.3 |
| | \$35,000+ | 57.9 | 50.1-65.8 |
| Insurance | Medicaid | 88.8 | 82.2-95.4 |
| | Insurance | 66.8 | 60.3-73.2 |

Stress cont.


**Prevalence of Individual Stressors
Tennessee, 2007**



**Prevalence of Stressors by Type*
Tennessee, 2007**




**Number of Stressor Types*
Tennessee, 2007**



* Analysis limited to women with at least one stressor.

Stress cont.

High vs. Low Stress Levels*
Tennessee, 2007




High Stress by Demographic and Socioeconomic Characteristics*
Tennessee, 2007

| | | Percent | 95% Confidence Interval |
|-------------------|----------------|---------|-------------------------|
| Race | White Non-Hisp | 34.5 | 28.1-40.9 |
| | Black Non-Hisp | 43.4 | 29.8-56.9 |
| Age (years) | <20 | 40.5 | 25.1-55.9 |
| | 20-29 | 38.2 | 30.8-45.6 |
| | 30+ | 25.2 | 15.5-34.9 |
| Married | Yes | 26.3 | 19.5-33.0 |
| | No | 44.3 | 35.5-53.0 |
| Education (years) | <12 | 37.6 | 25.8-49.4 |
| | 12 | 45.1 | 34.2-56.0 |
| | >12 | 26.7 | 19.2-34.2 |
| Household Income | <\$10,000 | 49.0 | 37.7-60.3 |
| | \$10-24,999 | 48.8 | 34.7-62.9 |
| | \$25-34,999 | 38.1 | 26.1-50.2 |
| | \$35,000+ | 11.6 | 4.7-18.5 |
| Insurance | Medicaid | 54.2 | 42.5-65.8 |
| | Insurance | 20.1 | 12.9-27.3 |


* Analysis limited to women with at least one stressor.

Stress cont.


Food Security*
Tennessee, 2007




Neighborhood Safety†
Tennessee, 2007



Racial Bias‡
Tennessee, 2007



Racial Bias by Maternal Race‡
Tennessee, 2007



* Food insecurity was defined as having eaten less than you felt you should in the 12 months prior to delivery because there wasn't enough money to buy food.

† An unsafe neighborhood was defined as one in which the mother always, often or sometimes felt unsafe in the 12 months prior to delivery.

‡ Racial bias was defined as having felt emotionally upset as a result of how you were treated based on your race in the 12 months prior to delivery.

Cigarette Smoking

Information regarding cigarette smoking was derived from the following questions:

Question #31: Have you smoked at least 100 cigarettes in the past 2 years?

- No
- Yes

Question #32: In the 3 months before you got pregnant, how many cigarettes did you smoke on an average day?

- 41 cigarettes or more
- 21 to 40 cigarettes
- 11 to 20 cigarettes
- 6 to 10 cigarettes
- 1 to 5 cigarettes
- Less than 1 cigarette
- None (0 cigarettes)


Additional (and similarly worded) questions about cigarette use in the last 3 months of pregnancy (Question #33) and cigarette use at the time of the survey (Question #34) may be found in the Appendix. Women who reported that they had not smoked at least 100 cigarettes in the past 2 years were classified as nonsmokers for all three time periods. Women with *any* reported cigarette use (even less than one cigarette) for a given time period were classified as smokers for that time period. Women who smoked prior to but not during pregnancy were classified as quitters. Quitters who reported smoking at the time of the survey were classified as having resumed smoking.

Key Findings:


- Approximately one-third of women (31.5%) smoked cigarettes in the 3 months prior to pregnancy.
- Among women who smoked prior to pregnancy, two-thirds continued to smoke during pregnancy, while one-third quit smoking.
- All women who were nonsmokers prior to pregnancy remained nonsmokers during pregnancy (i.e. no women initiated smoking while pregnant).
- Overall, one-fifth (21.2%) of women smoked during the last 3 months of pregnancy.
- White non-Hispanics were more likely than black non-Hispanics to smoke during pregnancy.
- Unmarried women were more likely than those who were married to smoke during pregnancy.
- Among women who quit smoking during pregnancy, approximately two-fifths had resumed smoking at the time of the survey.
- Women with unintended pregnancies were more likely than those with intended pregnancies to smoke prior to pregnancy. However, there was not a statistically significant difference in the prevalence of smoking in the last 3 months of pregnancy or in the percentage of women who quit.
- Women who received late or no prenatal care were more likely to smoke during pregnancy than those who received early prenatal care.
- Women who reported one or more stressors in the 12 months prior to delivery were more likely to smoke during pregnancy than those who did not report any stressors.

Cigarette Smoking cont.

**Cigarette Smoking During Pregnancy
Tennessee, 2007**



**Amount Smoked During Pregnancy*
Tennessee, 2007**




**Smoking During Pregnancy by Demographic and Socioeconomic Characteristics
Tennessee, 2007**

| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 26.9 | 21.7-32.1 |
| | Black Non-Hisp | 7.0 | 0.9-13.1 |
| | Hispanic | 8.1 | 0.0-18.5 |
| Age (years) | <20 | 30.2 | 17.1-43.2 |
| | 20-29 | 23.0 | 17.4-28.6 |
| | 30+ | 13.2 | 7.3-19.1 |
| Married | Yes | 15.2 | 10.8-19.6 |
| | No | 28.8 | 21.5-36.1 |
| Education (years) | <12 | 34.9 | 24.8-45.0 |
| | 12 | 28.6 | 19.7-37.4 |
| | >12 | 10.0 | 6.0-14.1 |
| Household Income | <\$10,000 | 29.7 | 20.3-39.2 |
| | \$10-24,999 | 34.4 | 22.1-46.7 |
| | \$25-34,999 | 19.8 | 10.9-28.8 |
| | \$35,000+ | 10.3 | 5.3-15.3 |
| Insurance | Medicaid | 30.1 | 20.4-39.8 |
| | Insurance | 10.6 | 6.3-14.9 |


* Analysis limited to women who smoked cigarettes in the last 3 months of pregnancy.

Cigarette Smoking *cont.*


**Smoking Prior to vs. During Pregnancy -- All Respondents
Tennessee, 2007**



**Smoking Prior to vs. During Pregnancy -- Prior Smokers*
Tennessee, 2007**



**Smoking During vs. After Pregnancy -- Quitters†
Tennessee, 2007**




* Analysis limited to women who smoked cigarettes in the 3 months prior to pregnancy.


† Analysis limited to women who quit smoking during pregnancy.

Cigarette Smoking cont.


**Cigarette Use by Pregnancy Intent
Tennessee, 2007**




**Smoking During Pregnancy by Prenatal Care Timing
Tennessee, 2007**



Quitting by Prenatal Care Timing
Tennessee, 2007**



**Smoking During Pregnancy by Presence of Stressors
Tennessee, 2007**



* Not statistically significant.

† Analysis limited to women who smoked cigarettes in the 3 months prior to pregnancy.

‡ Results for women with late or no care are based on an unweighted sample size of 50. Therefore, results may be unreliable and should be interpreted with caution.

Alcohol Use

Information regarding alcohol use was derived from the following questions:

Question #35: Have you had any alcoholic drinks in the past 2 years?

- No
- Yes

Question #36a: During the 3 months before you got pregnant, how many alcoholic drinks did you have in an average week?

- 14 drinks or more
- 7 to 13 drinks
- 4 to 6 drinks
- 1 to 3 drinks
- Less than 1 drink
- I didn't drink then

Question #36b: During the 3 months before you got pregnant, how many times did you drink 5 alcoholic drinks or more in one sitting?

- 6 or more times
- 4 to 5 times
- 2 to 3 times
- 1 time
- I didn't have 5 or more drinks in 1 sitting
- I didn't drink then

Additional (and similarly worded) questions about alcohol use and binge drinking in the last 3 months of pregnancy (Questions #37a and #37b) may be found in the Appendix. Women who reported not having any alcoholic drinks in the past 2 years were classified as nondrinkers for both time periods. Women with *any* reported alcohol use (even less than one drink) for a given time period were classified as drinkers for that time period, while those who drank 7 or more drinks per week were classified as frequent drinkers. Women who drank more than 5 drinks at a sitting at least once in a given time period were classified as binge drinkers for that time period. Women who drank prior to but not during pregnancy were classified as quitters.


Key Findings:

- Approximately two-fifths of women (42.8%) drank alcohol in the 3 months prior to pregnancy.
- Among women who drank prior to pregnancy, approximately 14% continued to do so during pregnancy, while the remaining 86% quit drinking.
- Overall, 5.8% of women drank alcohol during the last 3 months of pregnancy.*
- Among women who drank prior to pregnancy 7% did so frequently, while among those who drank during pregnancy there were no frequent drinkers.
- Among women who drank prior to pregnancy, approximately two-thirds engaged in binge drinking at least once, while among those who drank during pregnancy 6% were binge drinkers.


* Due to the small number of women who drank during pregnancy, it was not possible to analyze alcohol use during pregnancy by demographic, socioeconomic or other characteristics.

Alcohol Use *cont.*


**Alcohol Use Prior to Pregnancy
Tennessee, 2007**



**Alcohol Use During Pregnancy
Tennessee, 2007**




**Alcohol Use Prior to vs. During Pregnancy -- Prior Drinkers*
Tennessee, 2007**




* Analysis limited to women who drank alcohol in the 3 months prior to pregnancy.

Alcohol Use *cont.*


Amount of Alcohol Consumed Prior to Pregnancy*
Tennessee, 2007




Amount of Alcohol Consumed During Pregnancy†‡
Tennessee, 2007



Binge Drinking Prior to Pregnancy*
Tennessee, 2007



Binge Drinking During Pregnancy†‡
Tennessee, 2007



* Analysis limited to women who drank alcohol in the 3 months prior to pregnancy.

† Analysis limited to women who drank alcohol in the last 3 months of pregnancy.

‡ Results are based on an unweighted sample size of 44. Therefore, results may be unreliable and should be interpreted with caution.

Maternal Infections

Information regarding maternal infections was derived from the following questions:

Question #74: During your most recent pregnancy, did a doctor, nurse or other health care worker tell you that you had a urinary tract infection (UTI), a sexually transmitted disease (STD) or any vaginal infection, including bacterial vaginosis or Group B Strep (Beta Strep)?

- _No
- _Yes


Question #75: What disease or infection were you told you had?

- _Genital warts (HPV)
- _Herpes
- _Chlamydia
- _Gonorrhea
- _Pelvic inflammatory disease (PID)
- _Syphilis
- _Group B Strep (Beta Strep)
- _Bacterial vaginosis
- _Trichomoniasis (Trich)
- _Yeast infection
- _Urinary tract infection (UTI)
- _Other

Key Findings:

- Approximately two-fifths of women (40.2%) were diagnosed with a urinary tract infection, sexually transmitted disease or other vaginal infection during pregnancy.
- Teenage mothers were more likely than older mothers to have been diagnosed with an infection.
- There was not a statistically significant difference in the prevalence of maternal infections among black non-Hispanics versus white non-Hispanics.
- Among women who reported being diagnosed with an infection, urinary tract infections were the most frequently reported type of infection.
- Among women who reported being diagnosed with an infection, less than five percent reported being diagnosed with genital warts, herpes, trichomoniasis or gonorrhea.
- Women who had been physically abused prior to and/or during pregnancy were more likely than those who were not abused to have been diagnosed with an infection.
- Women who reported one or more stressors in the 12 months prior to delivery more likely than those who reported no stressors to have been diagnosed with an infection.

Maternal Infections *cont.*




**Maternal Infections by Demographic and Socioeconomic Characteristics
Tennessee, 2007**


| | | Percent <i>with</i> an Infection | 95% Confidence Interval |
|--------------------------|-----------------------|----------------------------------|-------------------------|
| Race | White Non-Hisp | 41.4 | 35.7-47.0 |
| | Black Non-Hisp | 36.9 | 24.9-48.8 |
| | Hispanic | 37.2 | 18.3-56.1 |
| Age (years) | <20 | 60.5 | 46.4-74.6 |
| | 20-29 | 39.7 | 33.2-46.2 |
| | 30+ | 30.9 | 22.7-39.0 |
| Married | Yes | 38.9 | 32.9-45.0 |
| | No | 41.8 | 33.7-49.8 |
| Education (years) | <12 | 45.9 | 35.3-56.5 |
| | 12 | 40.0 | 30.2-49.7 |
| | >12 | 36.8 | 30.2-43.4 |
| Household Income | <\$10,000 | 46.4 | 35.9-56.8 |
| | \$10-24,999 | 34.6 | 22.3-46.8 |
| | \$25-34,999 | 47.2 | 35.7-58.7 |
| | \$35,000+ | 33.6 | 26.1-41.0 |
| Insurance | Medicaid | 52.1 | 41.2-62.9 |
| | Insurance | 32.1 | 25.6-38.6 |

Maternal Infections *cont.*


**Maternal Infections by Type^{*†‡}
Tennessee, 2007**



**Maternal Infections by Presence of Stressors
Tennessee, 2007**



**Maternal Infections by Physical Abuse
Tennessee, 2007**



* Analysis limited to women with a diagnosed UTI, STD or other vaginal infection.

† Women may have reported more than one type of infection. Therefore, percentages do not sum to 100%.

‡ No women reported being diagnosed with PID or syphilis.

Maternal Health Problems

Information regarding maternal health problems was derived from the following questions:

Question #29: Did you have any of these problems during your most recent pregnancy?

- a. High blood sugar (diabetes) that started before this pregnancy
- b. High blood sugar (diabetes) that started during this pregnancy
- c. Vaginal bleeding
- d. Kidney or bladder (urinary tract) infection
- e. Severe nausea, vomiting or dehydration
- f. Cervix had to be sewn shut (incompetent cervix)
- g. High blood pressure, hypertension (including pregnancy-induced, preeclampsia or toxemia)
- h. Problems with the placenta (such as abruptio placentae or placenta previa)
- i. Labor pains more than 3 weeks before my baby was due (preterm or early labor)
- j. Water broke more than 3 weeks before my baby was due (premature rupture of membranes)
- k. I had to have a blood transfusion
- l. I was hurt in a car accident

Question #30: Did you do any of the following things because of these problems?

- a. I went to the hospital or emergency room and stayed less than one day
- b. I went to the hospital and stayed 1 to 7 days
- c. I went to the hospital and stayed more than 7 days
- d. I stayed in bed at home more than 2 days because of my doctor's or nurse's advise


Mothers who responded negatively to all of the items in Question #29 were classified as having no health problems during pregnancy. Those who responded positively to any of these items were classified as having health problems.

Key Findings:


- Approximately three-quarters (74.2%) of mothers reported at least one health problem during pregnancy.
- Among women with health problems, approximately 40% reported just one problem, while 12% reported 4 or more problems.
- Women on Medicaid or TennCare were more likely to have health problems than those with health insurance. The prevalence of maternal health problems was similar across other demographic and socioeconomic groups.
- The most frequently reported health problem was severe nausea, vomiting or dehydration.
- Approximately one-fifth of women reported a hospital stay of one day or greater because of their health problem(s).
- Approximately one-third of women reported bed rest of more than two days duration because of their health problem(s).

Maternal Health Problems *cont.*

**Health Problems During Pregnancy
Tennessee, 2007**



**Number of Health Problems*
Tennessee, 2007**




**Health Problems by Demographic and Socioeconomic Characteristics
Tennessee, 2007**

| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 74.9 | 69.8-79.9 |
| | Black Non-Hisp | 72.9 | 61.6-84.1 |
| | Hispanic | 64.7 | 45.5-84.0 |
| Age (years) | <20 | 80.0 | 68.4-91.8 |
| | 20-29 | 73.2 | 67.1-79.4 |
| | 30+ | 73.2 | 65.6-80.8 |
| Married | Yes | 75.1 | 69.8-80.4 |
| | No | 73.1 | 65.5-80.6 |
| Education (years) | <12 | 78.8 | 70.0-87.6 |
| | 12 | 78.3 | 69.8-86.9 |
| | >12 | 70.1 | 63.7-76.4 |
| Household Income | <\$10,000 | 78.6 | 69.7-87.6 |
| | \$10-24,999 | 61.9 | 48.8-74.9 |
| | \$25-34,999 | 79.7 | 70.5-88.9 |
| | \$35,000+ | 72.0 | 64.8-79.2 |
| Insurance | Medicaid | 81.5 | 72.6-90.3 |
| | Insurance | 68.0 | 61.3-74.6 |


* Analysis limited to women with one or more health problems.

Maternal Health Problems *cont.*


**Health Problems by Type^{*†}
Tennessee, 2007**



**Length of Hospital Stays for Health Problems^{*‡}
Tennessee, 2007**



**Bed Rest for Health Problems^{*}
Tennessee, 2007**



* Analysis limited to women with one or more health problems.

† Women may have reported more than one health problem. Therefore, percentages do not sum to 100%.

‡ Women may have had more than one hospital visit or stay. Therefore, categories are not mutually exclusive and percentages do not sum to 100%.

WIC Participation

Information regarding WIC participation was derived from the following question:

Question #28: During your most recent pregnancy, were you on WIC (the Supplemental Nutrition Program for Women, Infants and Children)?

_No

_Yes


Key Findings:

- Approximately one-half (50.9%) of women were enrolled in WIC during pregnancy.
- WIC participation increased with decreasing age. Almost all teenage mothers were enrolled in the program.
- Unmarried women were more likely to participate in WIC than married women.
- The prevalence of WIC participation was higher among women who reported food insecurity than among those not reporting food insecurity.*
- Approximately one-fourth (28.6%) of women who reported food insecurity were not enrolled in WIC. However, it was not possible to determine whether or not these women would have qualified for the program.
- WIC participants were less likely than non-participants to initiate breastfeeding.
- The most frequently reported reason for not breastfeeding among WIC participants was that they did not like breastfeeding.


* Food insecurity was defined as having eaten less than you felt you should in the 12 months prior to delivery because there wasn't enough money to buy food.

WIC Participation *cont.*

WIC Participation during Pregnancy
Tennessee, 2007



WIC Participation by Food Security Status
Tennessee, 2007




WIC Participation by Demographic and Socioeconomic Characteristics
Tennessee, 2007


| | | Percent | 95% Confidence Interval |
|-------------------|----------------|---------|-------------------------|
| Race | White Non-Hisp | 46.3 | 40.5-52.0 |
| | Black Non-Hisp | 60.5 | 48.4-72.6 |
| | Hispanic | 73.7 | 56.3-91.1 |
| Age (years) | <20 | 92.4 | 85.0-99.8 |
| | 20-29 | 51.1 | 44.4-57.8 |
| | 30+ | 30.1 | 22.0-38.1 |
| Married | Yes | 29.4 | 23.8-35.1 |
| | No | 78.3 | 71.4-85.1 |
| Education (years) | <12 | 74.3 | 64.9-83.9 |
| | 12 | 69.8 | 60.7-78.4 |
| | >12 | 27.2 | 20.8-33.6 |
| Household Income | <\$10,000 | 76.2 | 67.1-85.4 |
| | \$10-24,999 | 71.7 | 60.3-83.2 |
| | \$25-34,999 | 65.7 | 54.8-76.6 |
| | \$35,000+ | 8.3 | 3.8-12.8 |
| Insurance | Medicaid | 78.6 | 69.8-87.4 |
| | Insurance | 24.4 | 18.2-30.8 |

WIC Participation *cont.*

**Breastfeeding Initiation by WIC Participation
Tennessee, 2007**



**Reasons for Not Initiating Breastfeeding by WIC Participation[†]
Tennessee, 2007**



* Not statistically significant

[†] Women may have reported more than one reason for not breastfeeding. Therefore, percentages do not sum to 100%.

Breastfeeding

Information regarding breastfeeding was derived from the following questions:

Question #53: Did you ever breastfeed or pump breast milk to feed your new baby?

- No
- Yes

Question #55: Are you still breastfeeding or feeding pumped milk to your new baby?

- No
- Yes

Question #56: How many weeks or months did you breastfeed or pump milk to feed your baby?


- Weeks
- Months
- Less than 1 week

Additional questions addressing reasons for not initiating breastfeeding (Question #54) and for stopping breastfeeding (Question #57) may be found in the Appendix.

Key Findings:

- Approximately two-thirds (68.6%) of mothers initiated breastfeeding.
- Black non-Hispanics were less likely than white non-Hispanics to initiate breastfeeding.
- Teenage mothers were less likely than older mothers to initiate breastfeeding.
- Unmarried mothers were less likely than those who were married to initiate breastfeeding.
- Women with 12 or fewer years of education were less likely to initiate breastfeeding than those with higher levels of education.
- There was not a statistically significant difference in the percentage of women initiating breastfeeding among those who received early prenatal care versus those who received late/no care.
- There was not a statistically significant difference in the percentage of women initiating breastfeeding among those whose prenatal care included a discussion of breastfeeding versus those whose care did not include such a discussion.
- The most frequently reported reason for not initiating breastfeeding was not liking breastfeeding.
- Among women who initiated breastfeeding, approximately 5% breastfed for less than 1 week, while just over one-half breastfed for 9 weeks or longer.
- The most frequently reported reason for stopping breastfeeding was not producing enough milk.
- There was not a statistically significant difference in the percentage of women initiating breastfeeding among women with versus those without symptoms of postpartum depression (see page 58).

Breastfeeding *cont.*




**Breastfeeding Initiation by Demographic and Socioeconomic Characteristics
Tennessee, 2007**


| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 70.9 | 65.6-76.2 |
| | Black Non-Hisp | 50.4 | 37.8-63.0 |
| | Hispanic | 91.2 | 80.4-100.0 |
| Age (years) | <20 | 50.4 | 35.0-65.0 |
| | 20-29 | 70.6 | 64.3-77.0 |
| | 30+ | 73.6 | 65.7-81.5 |
| Married | Yes | 79.9 | 74.9-85.0 |
| | No | 54.3 | 46.0-62.6 |
| Education (years) | <12 | 58.6 | 47.9-69.4 |
| | 12 | 55.7 | 45.6-65.8 |
| | >12 | 81.7 | 76.4-86.9 |
| Household Income | <\$10,000 | 54.2 | 43.5-64.8 |
| | \$10-24,999 | 63.8 | 51.0-76.6 |
| | \$25-34,999 | 74.6 | 64.5-84.7 |
| | \$35,000+ | 79.6 | 73.0-86.2 |
| Insurance | Medicaid | 51.5 | 40.4-62.6 |
| | Insurance | 78.7 | 72.8-84.5 |

Breastfeeding *cont.*


Breastfeeding Initiation by Prenatal Care Timing*
Tennessee, 2007



Breastfeeding Initiation by Prenatal Care Content*
Tennessee, 2007



Reasons for Not Initiating Breastfeeding^{†‡}
Tennessee, 2007





* Not statistically significant.


† Women may have reported more than one reason for not breastfeeding. Therefore, percentages do not sum to 100%.

‡ Other reasons for not breastfeeding included "My milk didn't come in/I didn't produce enough milk/my milk was bad", "My baby wouldn't nurse", "My body was unhealthy", "I smoked", and "I just didn't want to".

Breastfeeding cont.



Reasons for Stopping Breastfeeding^{†‡} Tennessee, 2007



* All women who reported that they had only fed their baby breast milk (i.e. had not introduced formula, baby food, juice, cow's milk, water, sugar water or any other food) completed the survey when their babies were at least 10 weeks old. These women were included in the '9+ weeks' group.

† Women may have reported more than one reason for stopping breastfeeding. Therefore, percentages do not sum to 100%.

‡ Other reasons for stopping breastfeeding included "I did not produce enough milk/not enough milk for twins/milk dried up", "I was sick/I was taking medicine/I was tired", and "Baby didn't tolerate it/couldn't keep it down/wouldn't nurse".

Dental Care

Information regarding dental care was derived from the following questions:

Question #80: This question is about the care of your teeth during your most recent pregnancy. For each item, circle Y (yes) if it is true or circle N (no) if it is not true.

- a. I needed to see a dentist for a problem
- b. I went to a dentist or dental clinic
- c. A dental or other health care worker talked with me about how to care for my teeth and gums

Question #81: Have you ever had your teeth cleaned by a dentist or dental hygienist?

- No
 Yes

Question #82: When did you have your teeth cleaned by a dentist or dental hygienist? For each of the three time periods, circle Y (yes) if you had your teeth cleaned then or circle N (no) if you did not have your teeth cleaned then.


- a. Before my most recent pregnancy
- b. During my most recent pregnancy
- c. After my most recent pregnancy

Key Findings:


- Approximately one-third (28.9%) of mothers reported having a dental problem during pregnancy.
- Among women with a reported dental problem, over one-half did not go to a dentist or dental clinic during pregnancy.
- Overall, approximately two-thirds (66.6%) of women did not see a dentist during pregnancy.
- A majority of women (79.5%) who went to a dentist or dental clinic during pregnancy reported talking with a health care worker about the care of their teeth and gums.
- Approximately 6% of women had never had their teeth cleaned by a dentist or dental hygienist.
- Among women who had ever had their teeth cleaned, approximately one-third (34.1%) reported having a cleaning during their most recent pregnancy.
- Compared to women who received late or no prenatal care, those who received early care were more likely to:
 - Go to a dentist during pregnancy,
 - Go to a dentist for a dental problem, and
 - Have their teeth cleaned during pregnancy.

Dental Care cont.

Dental Care During Pregnancy*
Tennessee, 2007



Dental Problems and Dental Care
Tennessee, 2007




Did Not Go to Dentist by Demographic and Socioeconomic Characteristics*
Tennessee, 2007

| | | Percent | 95% Confidence Interval |
|-------------------|----------------|---------|-------------------------|
| Race | White Non-Hisp | 61.5 | 56.0-67.1 |
| | Black Non-Hisp | 76.1 | 65.9-86.3 |
| | Hispanic | 86.8 | 73.8-99.7 |
| Age (years) | <20 | 63.1 | 48.7-77.5 |
| | 20-29 | 72.1 | 66.2-77.9 |
| | 30+ | 57.6 | 49.0-66.1 |
| Married | Yes | 59.8 | 53.8-65.9 |
| | No | 75.2 | 68.1-82.2 |
| Education (years) | <12 | 74.6 | 65.1-84.1 |
| | 12 | 70.1 | 61.0-79.2 |
| | >12 | 60.3 | 53.8-66.9 |
| Household Income | <\$10,000 | 86.0 | 78.9-93.1 |
| | \$10-24,999 | 68.1 | 55.4-80.8 |
| | \$25-34,999 | 68.8 | 58.3-79.3 |
| | \$35,000+ | 46.9 | 39.0-54.8 |
| Insurance | Medicaid | 68.9 | 58.8-79.1 |
| | Insurance | 56.3 | 49.4-63.1 |


* Analysis includes all women regardless of whether they reported a dental problem.

Dental Care *cont.* †


**Discussion of Teeth and Gum Care
Tennessee, 2007**




**Teeth Ever Cleaned by Dentist
Tennessee, 2007**



**When Teeth Cleaned by Dentist†
Tennessee, 2007**



**Dental Care by Prenatal Care Timing
Tennessee, 2007**



* Analysis limited to women who had ever had their teeth cleaned by a dentist.

† Women may have had their teeth cleaned in one or more of the listed time periods. Therefore, percentages do not sum to 100%.

Postpartum Depression

Information regarding postpartum depression was derived from the following questions:

Question #79a: Since your new baby was born, how often have you felt down, depressed or hopeless?

- _Always
- _Often
- _Sometimes
- _Rarely
- _Never

Question #79b: Since your new baby was born, how often have you had little interest or little pleasure in doing things?


- _Always
- _Often
- _Sometimes
- _Rarely
- _Never

Women who responded ‘Always’ or ‘Often’ to one or both of the above questions were classified as having symptoms of postpartum depression.

Key Findings:

- Approximately 12% of mothers have often or always felt down, depressed, or hopeless since their baby was born, and 13% have often or always felt little interest or pleasure in doing things.
- Overall, approximately one-fifth (19.3%) of women reported symptoms of postpartum depression.
- Women with unintended pregnancies were more likely to have symptoms of postpartum depression than women with intended pregnancies.
- Women who were physically abused were more likely to have symptoms of postpartum depression than women who were not abused (see page 27).
- There was not a statistically significant difference in the prevalence of postpartum depression among women who reported one or more stressors versus those who did not. However, the percentage of women with symptoms of postpartum depression increased with increasing number of stressors.
- There was not a statistically significant difference in the percentage of women initiating breastfeeding among women with versus those without symptoms of postpartum depression.

Postpartum Depression *cont.*




Postpartum Depression by Demographic and Socioeconomic Characteristics Tennessee, 2007


| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 19.3 | 14.6-24.1 |
| | Black Non-Hisp | 20.5 | 10.6-30.4 |
| | Hispanic | 20.7 | 5.2-36.1 |
| Age (years) | <20 | 28.0 | 14.9-41.0 |
| | 20-29 | 19.9 | 14.4-25.5 |
| | 30+ | 13.8 | 7.8-19.8 |
| Married | Yes | 12.3 | 8.2-16.4 |
| | No | 28.2 | 20.7-35.6 |
| Education (years) | <12 | 26.9 | 17.0-36.8 |
| | 12 | 22.5 | 14.1-30.8 |
| | >12 | 13.9 | 9.1-18.7 |
| Household Income | <\$10,000 | 28.6 | 18.9-38.3 |
| | \$10-24,999 | 17.2 | 7.8-26.7 |
| | \$25-34,999 | 22.6 | 12.6-32.6 |
| | \$35,000+ | 8.3 | 4.0-12.7 |
| Insurance | Medicaid | 29.8 | 19.7-39.9 |
| | Insurance | 11.1 | 6.7-15.5 |

Postpartum Depression *cont.*


**Postpartum Depression by Pregnancy Intent
Tennessee, 2007**




**Postpartum Depression by Presence of Stressors*
Tennessee, 2007**



**Postpartum Depression by Number of Stressors†
Tennessee, 2007**



**Breastfeeding Initiation by Symptoms of Postpartum Depression*
Tennessee, 2007**



* Not statistically significant.

† Results for women with 6-13 stressors are based on an unweighted sample size of 58. Therefore, results may be unreliable and should be interpreted with caution.

Postpartum Birth Control

Information regarding postpartum birth control was derived from the following question:

Question #66: Are you or your husband or partner doing anything now to keep from getting pregnant?

No



Yes

Additional questions addressing reasons for not using birth control (Question #67) and birth control methods (Question #68) may be found in the Appendix.

Key Findings:

- The majority of mothers (89.2%) were using postpartum birth control at the time of the survey.
 - Use of postpartum birth control was similar across demographic and socioeconomic subgroups.
 - Among women who were using birth control, the most frequently used methods were condoms and birth control pills.
 - Among women who were not using birth control, the most frequently reported reason for not doing so was not wanting to.
- Among women who were *not* trying to get pregnant at the time of their most recent pregnancy, birth control use increased from approximately 52% prior to pregnancy to 91% postpartum.
- There was no statistically significant difference in the percentage of women using postpartum birth control among those with late or no prenatal care versus those with early care.
- There were no statistically significant differences in the percentage of women using postpartum birth control by prenatal care timing, prenatal care content, post partum checkup status, or breastfeeding status (data not shown).

Postpartum Birth Control *cont.*




Postpartum Birth Control by Demographic and Socioeconomic Characteristics Tennessee, 2007

| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 89.6 | 86.2-93.1 |
| | Black Non-Hisp | 90.2 | 83.2-97.2 |
| | Hispanic | 81.3 | 65.3-97.2 |
| Age (years) | <20 | 84.5 | 74.3-94.7 |
| | 20-29 | 90.3 | 86.3-94.2 |
| | 30+ | 89.3 | 84.3-94.4 |
| Married | Yes | 88.7 | 84.9-92.6 |
| | No | 89.7 | 84.8-94.6 |
| Education (years) | <12 | 85.3 | 78.0-92.6 |
| | 12 | 88.6 | 82.3-94.8 |
| | >12 | 91.5 | 87.8-95.1 |
| Household Income | <\$10,000 | 88.2 | 81.7-94.6 |
| | \$10-24,999 | 91.9 | 84.6-99.3 |
| | \$25-34,999 | 89.5 | 82.1-96.8 |
| | \$35,000+ | 90.2 | 85.6-94.8 |
| Insurance | Medicaid | 89.9 | 83.8-96.1 |
| | Insurance | 88.6 | 84.2-92.9 |


*Analysis limited to women not trying to get pregnant prior to most recent pregnancy, regardless of pregnancy intention.

Postpartum Birth Control *cont.*

**Postpartum Birth Control Methods^{*†}
Tennessee, 2007**



**Reasons for Not Using Postpartum Birth Control*
Tennessee, 2007**



* Women may have reported more than one birth control method or reason for not using birth control. Therefore, percentages do not sum to 100%.

† Diaphragm, contraceptive patch, and monthly shot are included in the 'other' category. Less than 1% of women reported using each of these birth control methods.

Postpartum Checkup

Information regarding postpartum checkups was derived from the following question:

Question #69: Since your new baby was born, have you had a postpartum checkup for yourself?

No


Yes

Key Findings:


- The majority of mothers (87.2%) had received a postpartum checkup for themselves.
- Married women were more likely than unmarried women to have had a postpartum checkup.
- Women with health insurance were more likely to have had a postpartum checkup than those on Medicaid or TennCare.
- Women who received early prenatal care were more likely to have had a postpartum checkup than those who received late or no care.
- There were no statistically significant differences in the percentage of women who had a postpartum checkup by pregnancy intent, breastfeeding status or symptoms of postpartum depression (data not shown).

Postpartum Checkup cont.

**Postpartum Checkup
Tennessee, 2007**



**Postpartum Checkup by Prenatal Care Timing
Tennessee, 2007**



**Postpartum Checkup by Demographic and Socioeconomic Characteristics
Tennessee, 2007**

| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 86.5 | 82.3-90.6 |
| | Black Non-Hisp | 89.0 | 80.8-97.2 |
| | Hispanic | 85.5 | 72.5-98.5 |
| Age (years) | <20 | 82.7 | 72.1-93.3 |
| | 20-29 | 87.1 | 82.5-91.7 |
| | 30+ | 89.7 | 83.7-95.7 |
| Married | Yes | 91.2 | 87.6-94.8 |
| | No | 82.3 | 76.0-88.6 |
| Education (years) | <12 | 79.1 | 70.2-88.0 |
| | 12 | 88.4 | 81.8-94.9 |
| | >12 | 90.8 | 86.6-94.9 |
| Household Income | <\$10,000 | 78.9 | 70.2-87.6 |
| | \$10-24,999 | 85.7 | 76.0-95.3 |
| | \$25-34,999 | 83.8 | 75.5-92.1 |
| | \$35,000+ | 95.5 | 92.2-98.8 |
| Insurance | Medicaid | 81.5 | 72.8-90.2 |
| | Insurance | 95.3 | 92.6-98.1 |

Infant Health Care

Information regarding infant health care was derived from the following questions:

Question #49: After your baby was born, was he or she put in an intensive care unit?

- No
- Yes
- I don't know

Question #50: After your baby was born, how long did he or she stay in the hospital?

- Less than 24 hours
- 24 to 48 hours
- 3 days
- 4 days
- 5 days
- 6 days or more
- My baby was not born in a hospital
- My baby is still in the hospital

Question #62: Was your new baby seen by a doctor, nurse or other health care worker during the first week after he or she left the hospital?

- No
- Yes

Question #63: Has your new baby had a well-baby checkup?

- No
- Yes


Additional questions addressing health insurance (Questions #64-65) may be found in the Appendix.

Key Findings:


- Approximately 8% of infants were placed in an intensive care unit (ICU) after delivery.
- Approximately two-thirds (63.5%) of infants spent 2 days or less in the hospital after delivery, while 6% of infants spent 6 days or more.
- Infants who were placed in an ICU were more likely to spend 6 or more days in the hospital than infants who were not placed in an ICU.
- Most women (94.8%) reported having health insurance for their new baby.
- Over one-half of women reported that their baby was covered by TennCare.
- Most infants were seen by a health care worker within the first week after leaving the hospital (86.0%) and had a well-baby visit (97.5%).
- Over 90% of infants in all demographic and socioeconomic subgroups had a well-baby visit.

Infant Health Care *cont.*


**Neonatal Intensive Care Unit
Tennessee, 2007**




**Infant Length of Hospital Stay*
Tennessee, 2007**




**Infant Length of Stay by Admission to ICU*
Tennessee, 2007**



**Health Insurance for Infant
Tennessee, 2007**



**Type of Infant Health Insurance†
Tennessee, 2007**




* Analysis excludes infants not born in a hospital.


† Women may have reported more than one type of insurance. Therefore, percentages do not sum to 100%.

Infant Health Care *cont.*

Infant Exam within 1 Week of Discharge*
Tennessee, 2007



Any Well-Baby Exam Since Discharge*
Tennessee, 2007



Well-Baby Exam by Demographic and Socioeconomic Characteristics*
Tennessee, 2007

| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 97.2 | 95.2-99.2 |
| | Black Non-Hisp | 97.5 | 93.4-100.0 |
| | Hispanic | 99.7 | 99.0-100.0 |
| Age (years) | <20 | 97.6 | 93.9-100.0 |
| | 20-29 | 97.4 | 95.1-99.6 |
| | 30+ | 97.7 | 94.4-100.0 |
| Married | Yes | 97.0 | 94.6-99.4 |
| | No | 98.1 | 95.9-100.0 |
| Education (years) | <12 | 96.0 | 91.6-100.0 |
| | 12 | 97.1 | 93.4-100.0 |
| | >12 | 98.4 | 96.9-100.0 |
| Household Income | <\$10,000 | 97.0 | 93.3-100.0 |
| | \$10-24,999 | 92.9 | 86.0-99.8 |
| | \$25-34,999 | 97.2 | 93.5-100.0 |
| | \$35,000+ | 99.9 | 99.8-100.0 |
| Insurance | Medicaid | 97.8 | 94.7-100.0 |
| | Insurance | 99.4 | 98.5-100.0 |

* Analysis limited to mothers whose infants were still alive and living with them. Mothers whose Infants were still in the hospital at the time of the survey were excluded from the analysis.

Sleep Behaviors

Information regarding sleep behaviors was derived from the following questions:

Question #60: How do you most often lay your baby down to sleep now?

- On his or her side
- On his or her back
- On his or her stomach

Question #61: How often does your new baby sleep in the same bed with you or anyone else?

- Always
- Often
- Sometimes
- Rarely
- Never


Women who reported that their new baby always, often or sometimes slept in the same bed were classified as bed sharing with their infant, while those who reported that their baby rarely or never slept in the same bed were classified as not bed sharing.

Key Findings:

- Approximately one-third (37.5%) of mothers most often used the stomach or side sleeping positions, while the remaining two-thirds reported using the back position most frequently.*
- Use of the stomach or side sleeping positions was similar across demographic and socioeconomic subgroups.
- Approximately one-half (50.9%) of mothers reported bed sharing with their infant.
- Bed sharing was more common among black non-Hispanics than among white non-Hispanics.
- Women less than 30 years old were more likely to bed share than were older women.
- Bed sharing increased with decreasing education and income.
- There were no statistically significant differences in sleeping behaviors by pregnancy intent, prenatal care, infant birthweight, breastfeeding, maternal postpartum checkup or well-baby visit (data not shown).

* *Infants who sleep on their stomachs and sides are at greater risk for Sudden Infant Death Syndrome (SIDS) than infants who sleep on their backs.*

Sleep Behaviors *cont.*



Stomach/Side Sleeping by Demographic and Socioeconomic Characteristics^{*†}


Tennessee, 2007

| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 36.1 | 30.4-41.9 |
| | Black Non-Hisp | 48.2 | 33.8-62.6 |
| | Hispanic | 31.2 | 10.1-52.4 |
| Age (years) | <20 | 40.0 | 24.6-55.4 |
| | 20-29 | 38.2 | 31.2-45.1 |
| | 30+ | 35.0 | 26.5-43.6 |
| Married | Yes | 36.3 | 30.2-42.4 |
| | No | 39.2 | 30.2-48.2 |
| Education (years) | <12 | 38.3 | 26.5-50.1 |
| | 12 | 33.4 | 23.4-43.3 |
| | >12 | 38.7 | 31.8-45.5 |
| Household Income | <\$10,000 | 38.2 | 26.8-49.6 |
| | \$10-24,999 | 34.9 | 21.5-48.3 |
| | \$25-34,999 | 40.8 | 28.9-52.7 |
| | \$35,000+ | 36.7 | 28.9-44.5 |
| Insurance | Medicaid | 40.8 | 28.9-52.8 |
| | Insurance | 37.2 | 30.3-44.2 |

* Analysis limited to mothers whose infants were still alive and living with them.

† Women were asked to select the single most often used sleeping position. Approximately 10% of women chose multiple sleeping positions and were excluded from the analysis.

Sleep Behaviors *cont.*



**Bed Sharing by Demographic and Socioeconomic Characteristics*
Tennessee, 2007**

| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 41.2 | 35.4-47.0 |
| | Black Non-Hisp | 79.1 | 69.8-88.5 |
| | Hispanic | 65.1 | 45.2-84.9 |
| Age (years) | <20 | 67.1 | 53.6-80.6 |
| | 20-29 | 54.2 | 47.5-61.0 |
| | 30+ | 36.2 | 27.6-44.8 |
| Married | Yes | 37.9 | 31.8-44.1 |
| | No | 67.4 | 59.7-75.1 |
| Education (years) | <12 | 72.2 | 62.6-81.9 |
| | 12 | 55.7 | 45.7-65.7 |
| | >12 | 36.7 | 29.9-43.5 |
| Household Income | <\$10,000 | 66.2 | 56.2-76.1 |
| | \$10-24,999 | 52.3 | 39.1-65.6 |
| | \$25-34,999 | 53.7 | 41.9-65.4 |
| | \$35,000+ | 31.3 | 23.8-38.8 |
| Insurance | Medicaid | 61.8 | 51.2-72.4 |
| | Insurance | 39.2 | 32.2-46.2 |

* Analysis limited to mothers whose infants were still alive and living with them.

Infant Smoke Exposure

Information regarding infant smoke exposure was derived from the following question:

Question #59: About how many hours a day, on average, is your new baby in the same room with someone who is smoking?

[#] Hours

_ Less than 1 hour a day

_ My baby is never in the same room with someone who is smoking


Women who reported that their infant was in the same room as a smoker for *any* amount of time (even less than one hour a day) were classified as having an infant exposed to secondhand smoke.

Key Findings:


- Approximately 9% of mothers reported that their infants were exposed to secondhand smoke.
- Infant exposure to secondhand smoke was more common among white non-Hispanics than among black non-Hispanics.
- Unmarried women were more likely than married women to report infant exposure to secondhand smoke.
- Teenage mothers were more likely than older mothers to report infant exposure to secondhand smoke.
- Women who smoked prior to, during or after pregnancy were more likely to report that their infant was in the same room with someone who was smoking than were women who did not smoke during each of these three time periods.

Infant Smoke Exposure *cont.*

**Infant Exposure to Secondhand Smoke*
Tennessee, 2007**



**Infant Smoke Exposure by Maternal Cigarette Smoking*
Tennessee, 2007**



**Infant Smoke Exposure by Demographic and Socioeconomic Characteristics*
Tennessee, 2007**

| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 11.1 | 7.1-15.1 |
| | Black Non-Hisp | 4.1 | 0.0-8.7 |
| | Hispanic | 0.0 | -- |
| Age (years) | <20 | 19.3 | 7.9-30.7 |
| | 20-29 | 8.2 | 4.3-12.2 |
| | 30+ | 4.4 | 0.4-8.4 |
| Married | Yes | 4.2 | 1.6-6.7 |
| | No | 14.5 | 8.5-20.5 |
| Education (years) | <12 | 13.2 | 5.8-20.5 |
| | 12 | 14.1 | 6.5-21.7 |
| | >12 | 3.6 | 0.9-6.3 |
| Household Income | <\$10,000 | 13.0 | 5.3-20.7 |
| | \$10-24,999 | 19.0 | 8.4-29.6 |
| | \$25-34,999 | 6.2 | 0.8-11.5 |
| | \$35,000+ | 3.3 | 0.3-6.3 |
| Insurance | Medicaid | 4.6 | 1.6-7.6 |
| | Insurance | 13.9 | 5.9-21.9 |

* Analysis limited to mothers whose infants were still alive and living with them.

Infant Safety

Information regarding infant safety was derived from the following question:

Question #77: Listed below are some statements about safety. For each one, circle Y (yes) if it applies to you or circle N (no) if it does not.

- a. My infant was brought home from the hospital in an infant car seat
- b. My baby always or almost always rides in an infant car seat
- c. My home has a working smoke alarm
- d. There are loaded guns, rifles or other firearms in my home


Key Findings:

- Almost all mothers reported using a car seat to bring their infant home from the hospital (99.2%) and always or almost always using a car seat (99.7%).*
- Approximately 5% of women reported that there is not a working smoke alarm in their home.*
- Approximately 8% of women reported that there are loaded firearms in their home.*


* Due to the small number of women with infant safety risk factors, it was not possible to analyze these variables by demographic, socioeconomic or other characteristics.

Infant Safety *cont.*


Car Seat Use After Hospital Discharge*
Tennessee, 2007




Car Seat Use Always or Almost Always*
Tennessee, 2007



Working Smoke Alarm in Home*
Tennessee, 2007



Loaded Firearms in Home*
Tennessee, 2007



* Analysis limited to mothers whose infants were still alive and living with them.


Low Birthweight

Information regarding infant birthweight was derived from birth certificates. Infants with birthweights of less than 2,500 grams were classified as low birthweight, while those with birthweights of 2,500 grams or more were classified as normal birthweight.

Key Findings:

- Approximately 8.5% of mothers gave birth to low birthweight infants.
- Black non-Hispanics were more likely than white-non-Hispanics to have a low birthweight infant.
- Unmarried women were more likely than those who were married to have a low birthweight infant.
- Women on Medicaid or TennCare were more likely to have a low birthweight infant than those with health insurance.
- There were no statistically significant differences in the prevalence of low birthweight by pregnancy intent, prenatal care, physical abuse, stress, maternal smoking, maternal infection, or dental care (data not shown).

Low Birthweight *cont.*



**Low Birthweight by Demographic and Socioeconomic Characteristics
Tennessee, 2007**

| | | Percent | 95% Confidence Interval |
|--------------------------|-----------------------|---------|-------------------------|
| Race | White Non-Hisp | 7.2 | 6.5-7.9 |
| | Black Non-Hisp | 13.2 | 9.5-16.9 |
| | Hispanic | 6.7 | 2.9-10.5 |
| Age (years) | <20 | 6.8 | 4.0-9.6 |
| | 20-29 | 8.5 | 7.4-9.6 |
| | 30+ | 9.5 | 7.6-11.4 |
| Married | Yes | 7.2 | 6.4-8.0 |
| | No | 10.2 | 8.4-12.0 |
| Education (years) | <12 | 7.6 | 5.5-9.6 |
| | 12 | 10.7 | 8.3-13.1 |
| | >12 | 7.9 | 6.8-9.0 |
| Household Income | <\$10,000 | 10.1 | 7.7-12.6 |
| | \$10-24,999 | 5.7 | 3.5-8.0 |
| | \$25-34,999 | 7.3 | 5.1-9.5 |
| | \$35,000+ | 8.4 | 6.9-9.9 |
| Insurance | Medicaid | 11.7 | 8.8-14.6 |
| | Insurance | 7.7 | 6.6-8.9 |

Appendix – 2007 TN PRAMS Questionnaire

PRAMS

Tennessee Pregnancy Risk Assessment Monitoring System

*A Survey of the Health of
Mothers and Babies
in Tennessee*

*For further information,
please call toll-free
1-877-984-8662*




Tennessee Department of Health
Office of Policy, Planning and Assessment
425 5th Avenue North, 4th Floor
Nashville, Tennessee 37243



First, we would like to ask a few questions about you and the time before you got pregnant with your new baby. Please check the box next to your answer.

1. *Just before you got pregnant, did you have health insurance?* Do not count Medicaid, TennCare or CoverKids.

- No
 Yes

2. *Just before you got pregnant, were you on Medicaid, TennCare or CoverKids?*

- No
 Yes

3. *During the month before you got pregnant with your new baby, how many times a week did you take a multivitamin or a prenatal vitamin?* These are pills that contain many different vitamins and minerals.

- I didn't take a multivitamin or a prenatal vitamin at all
 1 to 3 times a week
 4 to 6 times a week
 Every day of the week

4. What is *your* date of birth?

19
 Month Day Year

5. *Just before you got pregnant with your new baby, how much did you weigh?*

Pounds OR Kilos

6. How tall are you without shoes?

Feet Inches

OR Centimeters

7. Would you say that, in general, your health is—

- Excellent
 Very good
 Good
 Fair
 Poor

8. *Before you got pregnant with your new baby, did you talk with a doctor, nurse, or other health care worker to prepare for a healthy pregnancy and baby?*

- No
 Yes

9. *Before you got pregnant with your new baby, did you ever have any other babies who were born alive?*

- No → **Go to Page 2, Question 12**
 Yes

10. Did the baby born *just before* your new one weigh 5 pounds, 8 ounces (2.5 kilos) or less at birth?

- No
 Yes

11. Was the baby *just before* your new one born more than 3 weeks before its due date?

- No
 Yes

2

The next questions are about the time when you got pregnant with your *new* baby.

12. Thinking back to just before you got pregnant with your *new* baby, how did you feel about becoming pregnant?

Check one answer

- I wanted to be pregnant sooner
- I wanted to be pregnant later
- I wanted to be pregnant then
- I didn't want to be pregnant then or at any time in the future

If you wanted to be pregnant later, answer Question 13. Otherwise, go to Question 14.

13. How much later did you want to become pregnant?

- Less than 1 year
- 1 year to less than 2 years
- 2 years to less than 3 years
- 3 years to less than 4 years
- 4 years or more

14. When you got pregnant with your new baby, were you trying to get pregnant?

- No
- Yes → Go to Question 18

15. When you got pregnant with your new baby, were you or your husband or partner doing anything to keep from getting pregnant?

(Some things people do to keep from getting pregnant include not having sex at certain times [rhythm] or withdrawal, and using birth control methods such as the pill, condoms, cervical ring, IUD, having their tubes tied, or their partner having a vasectomy.)

- No
- Yes → Go to Question 17

16. What were your or your husband's or partner's reasons for not doing anything to keep from getting pregnant?

Check all that apply

- I didn't mind if I got pregnant
- I thought I could not get pregnant at that time
- I had side effects from the birth control method I was using
- I had problems getting birth control when I needed it
- I thought my husband or partner or I was sterile (could not get pregnant at all)
- My husband or partner didn't want to use anything
- Other → Please tell us:

If you were not doing anything to keep from getting pregnant, go to Question 18.

17. When you got pregnant with your new baby, what were you or your husband or partner doing to keep from getting pregnant?

Check all that apply

- Tubes tied or closed (female sterilization)
- Vasectomy (male sterilization)
- Pill
- Condoms
- Shot once a month (Lunelle®)
- Shot once every 3 months (Depo-Provera®)
- Contraceptive patch (OrthoEvra®)
- Diaphragm, cervical cap, or sponge
- Cervical ring (NuvaRing® or others)
- IUD (including Mirena®)
- Rhythm method or natural family planning
- Withdrawal (pulling out)
- Not having sex (abstinence)
- Other _____ → Please tell us:

The next questions are about the prenatal care you received during your most recent pregnancy. Prenatal care includes visits to a doctor, nurse, or other health care worker before your baby was born to get checkups and advice about pregnancy. (It may help to look at the calendar when you answer these questions.)

18. How many weeks or months pregnant were you when you were *sure* you were pregnant? (For example, you had a pregnancy test or a doctor or nurse said you were pregnant.)

____ Weeks OR ____ Months

- I don't remember

19. How many weeks or months pregnant were you when you had your first visit for prenatal care? Do not count a visit that was only for a pregnancy test or only for WIC (the Special Supplemental Nutrition Program for Women, Infants, and Children).

____ Weeks OR ____ Months

- I didn't go for prenatal care

4

20. Did you get prenatal care as early in your pregnancy as you wanted?

- No
- Yes
- I didn't want prenatal care

→ **Go to Question 22**

21. Here is a list of problems some women can have getting prenatal care. For each item, circle **Y** (Yes) if it was a problem for you during your most recent pregnancy or circle **N** (No) if it was not a problem or did not apply to you.

| | No | Yes |
|---|----|-----|
| a. I couldn't get an appointment when I wanted one | N | Y |
| b. I didn't have enough money or insurance to pay for my visits | N | Y |
| c. I had no way to get to the clinic or doctor's office | N | Y |
| d. I couldn't take time off from work | N | Y |
| e. The doctor or my health plan would not start care as early as I wanted | N | Y |
| f. I didn't have my Medicaid/TennCare/CoverKids card | N | Y |
| g. I had no one to take care of my children | N | Y |
| h. I had too many other things going on | N | Y |
| i. I didn't want anyone to know I was pregnant | N | Y |
| j. Other | N | Y |

Please tell us:

If you did not go for prenatal care, go to Question 24.

22. How was your prenatal care paid for?

Check all that apply

- Medicaid
- Personal income (cash, check, or credit card)
- Health insurance or HMO (including insurance from your work or your husband's work)
- TennCare
- CoverKids
- Other → Please tell us:

23. During any of your prenatal care visits, did a doctor, nurse, or other health care worker talk with you about any of the things listed below? Please count only discussions, not reading materials or videos. For each item, circle **Y** (Yes) if someone talked with you about it or circle **N** (No) if no one talked with you about it.

| | No | Yes |
|--|----|-----|
| a. How smoking during pregnancy could affect my baby | N | Y |
| b. Breastfeeding my baby | N | Y |
| c. How drinking alcohol during pregnancy could affect my baby | N | Y |
| d. Using a seat belt during my pregnancy | N | Y |
| e. Birth control methods to use after my pregnancy | N | Y |
| f. Medicines that are safe to take during my pregnancy | N | Y |
| g. How using illegal drugs could affect my baby | N | Y |
| h. Doing tests to screen for birth defects or diseases that run in my family | N | Y |
| i. What to do if my labor starts early | N | Y |
| j. Getting tested for HIV (the virus that causes AIDS) | N | Y |
| k. Physical abuse to women by their husbands or partners | N | Y |

24. At any time during your most recent pregnancy or delivery, did you have a test for HIV (the virus that causes AIDS)?

- No
- Yes → **Go to Question 26**
- I don't know

25. Were you offered an HIV test during your most recent pregnancy or delivery?

- No
- Yes

26. Have you ever heard or read that taking the vitamin folic acid can help prevent some birth defects?

- No → **Go to Question 28**
- Yes

27. Have you ever heard about folic acid from any of the following?

Check all that apply

- Magazine or newspaper article
- Radio or television
- Doctor, nurse, or other health care worker
- Book
- Family or friends
- Other → Please tell us:

The next questions are about your most recent pregnancy and things that might have happened during your pregnancy.

28. During your most recent pregnancy, were you on WIC (the Special Supplemental Nutrition Program for Women, Infants, and Children)?

- No
- Yes

29. Did you have any of these problems during your most recent pregnancy? For each item, circle Y (Yes) if you had the problem or circle N (No) if you did not.

| | No | Yes |
|---|----|-----|
| a. High blood sugar (diabetes) that started <i>before</i> this pregnancy | N | Y |
| b. High blood sugar (diabetes) that started <i>during</i> this pregnancy | N | Y |
| c. Vaginal bleeding | N | Y |
| d. Kidney or bladder (urinary tract) infection | N | Y |
| e. Severe nausea, vomiting, or dehydration | N | Y |
| f. Cervix had to be sewn shut (incompetent cervix) | N | Y |
| g. High blood pressure, hypertension (including pregnancy-induced hypertension [PIH]), preeclampsia, or toxemia | N | Y |
| h. Problems with the placenta (such as abruptio placentae or placenta previa) | N | Y |
| i. Labor pains more than 3 weeks before my baby was due (preterm or early labor) | N | Y |
| j. Water broke more than 3 weeks before my baby was due (premature rupture of membranes [PROM]) | N | Y |
| k. I had to have a blood transfusion | N | Y |
| l. I was hurt in a car accident | N | Y |

If you did not have any of these problems, go to Page 6, Question 31.

6

30. Did you do any of the following things because of these problems? For each item, circle **Y** (Yes) if you did that thing or circle **N** (No) if you did not.

| | No | Yes |
|--|----|-----|
| a. I went to the hospital or emergency room and stayed less than 1 day | N | Y |
| b. I went to the hospital and stayed 1 to 7 days | N | Y |
| c. I went to the hospital and stayed more than 7 days | N | Y |
| d. I stayed in bed at home more than 2 days because of my doctor's or nurse's advice | N | Y |

The next questions are about smoking cigarettes and drinking alcohol.

31. Have you smoked at least 100 cigarettes in the past 2 years? (A pack has 20 cigarettes.)

- No → Go to Question 35
 Yes

32. In the 3 months before you got pregnant, how many cigarettes did you smoke on an average day? (A pack has 20 cigarettes.)

- 41 cigarettes or more
- 21 to 40 cigarettes
- 11 to 20 cigarettes
- 6 to 10 cigarettes
- 1 to 5 cigarettes
- Less than 1 cigarette
- None (0 cigarettes)

33. In the last 3 months of your pregnancy, how many cigarettes did you smoke on an average day? (A pack has 20 cigarettes.)

- 41 cigarettes or more
- 21 to 40 cigarettes
- 11 to 20 cigarettes
- 6 to 10 cigarettes
- 1 to 5 cigarettes
- Less than 1 cigarette
- None (0 cigarettes)

34. How many cigarettes do you smoke on an average day now? (A pack has 20 cigarettes.)

- 41 cigarettes or more
- 21 to 40 cigarettes
- 11 to 20 cigarettes
- 6 to 10 cigarettes
- 1 to 5 cigarettes
- Less than 1 cigarette
- None (0 cigarettes)

35. Have you had any alcoholic drinks in the past 2 years? (A drink is 1 glass of wine, wine cooler, can or bottle of beer, shot of liquor, or mixed drink.)

- No → Go to Question 38
 Yes

36a. During the 3 months before you got pregnant, how many alcoholic drinks did you have in an average week?

- 14 drinks or more a week
- 7 to 13 drinks a week
- 4 to 6 drinks a week
- 1 to 3 drinks a week
- Less than 1 drink a week
- I didn't drink then

36b. During the 3 months before you got pregnant, how many times did you drink 5 alcoholic drinks or more in one sitting?

- 6 or more times
- 4 to 5 times
- 2 to 3 times
- 1 time
- I didn't have 5 drinks or more in 1 sitting
- I didn't drink then

37a. During the last 3 months of your pregnancy, how many alcoholic drinks did you have in an average week?

- 14 drinks or more a week
- 7 to 13 drinks a week
- 4 to 6 drinks a week
- 1 to 3 drinks a week
- Less than 1 drink a week
- I didn't drink then

37b. During the last 3 months of your pregnancy, how many times did you drink 5 alcoholic drinks or more in one sitting?

- 6 or more times
- 4 to 5 times
- 2 to 3 times
- 1 time
- I didn't have 5 drinks or more in 1 sitting
- I didn't drink then

Pregnancy can be a difficult time for some women. The next questions are about things that may have happened before and during your most recent pregnancy.

38. This question is about things that may have happened during the 12 months before your new baby was born. For each item, circle **Y** (Yes) if it happened to you or circle **N** (No) if it did not. (It may help to use the calendar.)

| | No | Yes |
|--|----|-----|
| a. A close family member was very sick and had to go into the hospital | N | Y |
| b. I got separated or divorced from my husband or partner | N | Y |
| c. I moved to a new address | N | Y |
| d. I was homeless | N | Y |
| e. My husband or partner lost his job | N | Y |
| f. I lost my job even though I wanted to go on working | N | Y |
| g. I argued with my husband or partner more than usual | N | Y |
| h. My husband or partner said he didn't want me to be pregnant | N | Y |
| i. I had a lot of bills I couldn't pay | N | Y |
| j. I was in a physical fight | N | Y |
| k. I or my husband or partner went to jail | N | Y |
| l. Someone very close to me had a bad problem with drinking or drugs | N | Y |
| m. Someone very close to me died | N | Y |

39. During the 12 months before your new baby was born, did you ever eat less than you felt you should because there wasn't enough money to buy food?

- No
- Yes

8

40. During the 12 months before your new baby was born, how often did you feel unsafe in the neighborhood where you lived?

- Always
- Often
- Sometimes
- Rarely
- Never

41. During the 12 months before your new baby was born, did you feel emotionally upset (for example angry, sad, or frustrated) as a result of how you were treated based on your race?

- No
- Yes

The next questions are about the time during the 12 months before you got pregnant with your new baby.

42a. During the 12 months before you got pregnant, did an ex-husband or ex-partner push, hit, slap, kick, choke, or physically hurt you in any other way?

- No
- Yes

42b. During the 12 months before you got pregnant, were you physically hurt in any way by your husband or partner?

- No
- Yes

The next questions are about the time during your most recent pregnancy.

43a. During your most recent pregnancy, did an ex-husband or ex-partner push, hit, slap, kick, choke, or physically hurt you in any other way?

- No
- Yes

43b. During your most recent pregnancy, were you physically hurt in any way by your husband or partner?

- No
- Yes

The next questions are about your labor and delivery. (It may help to look at the calendar when you answer these questions.)

44. When was your baby due?

Month Day Year

45. When did you go into the hospital to have your baby?

Month Day Year

I didn't have my baby in a hospital

46. When was your baby born?

Month Day Year

47. When were you discharged from the hospital after your baby was born? (It may help to use the calendar.)

 Month Day Year

I didn't have my baby in a hospital

48. How was your delivery paid for?

Check all that apply

- Medicaid
- Personal income (cash, check, or credit card)
- Health insurance or HMO (including insurance from your work or your husband's work)
- TennCare
- CoverKids
- Other → Please tell us:

The next questions are about the time since your new baby was born.

49. After your baby was born, was he or she put in an intensive care unit?

- No
- Yes
- I don't know

50. After your baby was born, how long did he or she stay in the hospital?

- Less than 24 hours (less than 1 day)
- 24 to 48 hours (1 to 2 days)
- 3 days
- 4 days
- 5 days
- 6 days or more
- My baby was not born in a hospital
- My baby is still in the hospital → **Go to Question 53**

51. Is your baby alive now?

- No → **Go to Page 11, Question 66**
- Yes

52. Is your baby living with you now?

- No → **Go to Page 11, Question 66**
- Yes

53. Did you ever breastfeed or pump breast milk to feed your new baby after delivery?

- No
- Yes → **Go to Page 10, Question 55**

10

54. What were your reasons for not breastfeeding your new baby?

Check all that apply

- My baby was sick and could not breastfeed
- I was sick or on medicine
- I had other children to take care of
- I had too many household duties
- I didn't like breastfeeding
- I didn't want to be tied down
- I was embarrassed to breastfeed
- I went back to work or school
- I wanted my body back to myself
- Other _____ → Please tell us:

If you did not breastfeed your new baby, go to Question 59.

55. Are you still breastfeeding or feeding pumped milk to your new baby?

No

Yes → **Go to Question 58**

56. How many weeks or months did you breastfeed or pump milk to feed your baby?

_____ Weeks **OR** _____ Months

Less than 1 week

57. What were your reasons for stopping breastfeeding?

Check all that apply

- My baby had difficulty nursing
- Breast milk alone did not satisfy my baby
- I thought my baby was not gaining enough weight
- My baby got sick and could not breastfeed
- My nipples were sore, cracked, or bleeding
- I thought I was not producing enough milk
- I had too many other household duties
- I felt it was the right time to stop breastfeeding
- I got sick and could not breastfeed
- I went back to work or school
- I wanted or needed someone else to feed the baby
- My baby was jaundiced (yellowing of the skin or whites of the eyes)
- Other _____ → Please tell us:

58. How old was your baby the first time you fed him or her anything besides breast milk? Include formula, baby food, juice, cow's milk, water, sugar water, or anything else you fed your baby.

_____ Weeks **OR** _____ Months

My baby was less than 1 week old

I have not fed my baby anything besides breast milk

If your baby is still in the hospital, go to Question 64.

59. About how many hours a day, on average, is your new baby in the same room with someone who is smoking?

_____ Hours

- Less than 1 hour a day
- My baby is never in the same room with someone who is smoking

60. How do you *most often* lay your baby down to sleep now?

Check one answer

- On his or her side
- On his or her back
- On his or her stomach

61. How often does your new baby sleep in the same bed with you or anyone else?

- Always
- Often
- Sometimes
- Rarely
- Never

62. Was your new baby seen by a doctor, nurse, or other health care worker during the first week after he or she left the hospital?

- No
- Yes

63. Has your new baby had a well-baby checkup?
(A well-baby checkup is a regular health visit for your baby, usually at 2, 4, or 6 months of age.)

- No
- Yes

64. Do you have health insurance, Medicaid, TennCare or CoverKids for your new baby?

- No → **Go to Question 66**
- Yes

65. What type of insurance is your new baby covered by?

Check all that apply

- Medicaid
- Private insurance or HMO (including insurance from your work or your husband's work)
- TennCare
- CoverKids
- Other → Please tell us:

66. Are you or your husband or partner doing anything *now* to keep from getting pregnant?

(Some things people do to keep from getting pregnant include not having sex at certain times [rhythm] or withdrawal, and using birth control methods such as the pill, condoms, cervical ring, IUD, having their tubes tied, or their partner having a vasectomy.)

- No
- Yes → **Go to Page 12, Question 68**

12

67. What are your or your husband's or partner's reasons for not doing anything to keep from getting pregnant now?

Check all that apply

- I am not having sex
- I want to get pregnant
- I don't want to use birth control
- My husband or partner doesn't want to use anything
- I don't think I can get pregnant (sterile)
- I can't pay for birth control
- I am pregnant now
- Other _____ → Please tell us:

If you are not doing anything to keep from getting pregnant now, go to Question 69.

68. What kind of birth control are you or your husband or partner using now to keep from getting pregnant?

Check all that apply

- Tubes tied or closed (female sterilization)
- Vasectomy (male sterilization)
- Pill
- Condoms
- Shot once a month (Lunelle®)
- Shot once every 3 months (Depo-Provera®)
- Contraceptive patch (OrthoEvra®)
- Diaphragm, cervical cap, or sponge
- Cervical ring (NuvaRing® or others)
- IUD (including Mirena®)
- Rhythm method or natural family planning
- Withdrawal (pulling out)
- Not having sex (abstinence)
- Other _____ → Please tell us:

69. Since your new baby was born, have you had a postpartum checkup for yourself?

(A postpartum checkup is the regular checkup a woman has after she gives birth.)

- No
- Yes

The next few questions are about the time during the 12 months before your new baby was born.

70. During the 12 months before your new baby was born, what were the sources of your household's income?

Check all that apply

- Paycheck or money from a job
- Money from family or friends
- Money from a business, fees, dividends, or rental income
- Aid such as Temporary Assistance for Needy Families (TANF), welfare, WIC, public assistance, general assistance, food stamps, or Supplemental Security Income
- Unemployment benefits
- Child support or alimony
- Social security, workers' compensation, disability, veteran benefits, or pensions
- Other _____ → Please tell us:

71. During the 12 months before your new baby was born, what was your total household income before taxes? Include your income, your husband's or partner's income, and any other income you may have used. (All information will be kept private and will not affect any services you are now getting.)

Check one answer

- Less than \$8,000
- \$8,000 to \$9,999
- \$10,000 to \$14,999
- \$15,000 to \$19,999
- \$20,000 to \$24,999
- \$25,000 to \$34,999
- \$35,000 to \$49,999
- \$50,000 or more

72. During the 12 months before your new baby was born, how many people, including yourself, depended on this income?

_____ People

The next few questions are on a variety of topics.

73. Which of the following statements best describes you during the 3 months before you got pregnant?

Check one answer

- I was trying to get pregnant
- I wasn't trying to get pregnant or trying to keep from getting pregnant
- I was trying to keep from getting pregnant but was not trying very hard
- I was trying hard to keep from getting pregnant

74. During your most recent pregnancy, did a doctor, nurse, or other health care worker tell you that you had a urinary tract infection (UTI), a sexually transmitted disease (STD), or any vaginal infection, including bacterial vaginosis or Group B Strep (Beta Strep)?

- No → Go to Question 76
- Yes

75. What disease or infection were you told you had?

Check all that apply

- Genital warts (HPV)
- Herpes
- Chlamydia
- Gonorrhea
- Pelvic inflammatory disease (PID)
- Syphilis
- Group B Strep (Beta Strep)
- Bacterial vaginosis
- Trichomoniasis (Trich)
- Yeast infections
- Urinary tract infection (UTI)
- Other → Please tell us:

76. Are you currently in school or working outside the home?

- No
- Yes

If your baby is not alive or is not living with you, go to Question 79a.

77. Listed below are some statements about safety. For each one, circle **Y** (Yes) if it applies to you or circle **N** (No) if it does not.

- | | No | Yes |
|---|----|-----|
| a. My infant was brought home from the hospital in an infant car seat | N | Y |
| b. My baby always or almost always rides in an infant car seat | N | Y |
| c. My home has a working smoke alarm | N | Y |
| d. There are loaded guns, rifles, or other firearms in my home | N | Y |

78. When your new baby rides in a car, truck, or van, how often does he or she ride in an infant car seat?

- Always
- Often
- Sometimes
- Rarely
- Never

79a. Since your new baby was born, how often have you felt down, depressed, or hopeless?

- Always
- Often
- Sometimes
- Rarely
- Never

79b. Since your new baby was born, how often have you had little interest or little pleasure in doing things?

- Always
- Often
- Sometimes
- Rarely
- Never

80. This question is about the care of your teeth during your most recent pregnancy. For each item, circle **Y** (Yes) if it is true or circle **N** (No) if it is not true.

- | | No | Yes |
|--|----|-----|
| a. I needed to see a dentist for a problem | N | Y |
| b. I went to a dentist or dental clinic | N | Y |
| c. A dental or other health care worker talked with me about how to care for my teeth and gums | N | Y |

81. Have you ever had your teeth cleaned by a dentist or dental hygienist?

- No → **Go to Question 83**
- Yes



82. When did you have your teeth cleaned by a dentist or dental hygienist? For each of the three time periods, circle **Y** (Yes) if you had your teeth cleaned then or circle **N** (No) if you did not have your teeth cleaned then.

- | | No | Yes |
|--|----|-----|
| a. Before my most recent pregnancy | N | Y |
| b. During my most recent pregnancy | N | Y |
| c. After my most recent pregnancy | N | Y |

83. What is today's date?

| | | |
|---|---|---|
| <input style="width: 100%;" type="text"/> | <input style="width: 100%;" type="text"/> | <input style="width: 100%;" type="text"/> |
| Month | Day | Year |

Please use this space for any additional comments you would like to make about the health of mothers and babies in Tennessee.



*Thanks for answering our questions!
Your answers will help us work to make Tennessee
mothers and babies healthier.*

December 7, 2006

